


ISSN-0011-7269

THE DECCAN GEOGRAPHER

Volume 61 Number 1 June and Number 2 December 2023

(UGC Care Listed: A Peer Review and Refereed Geography Journal)



Chief Editor
Professor B. C. Vaidya

THE DECCAN GEOGRAPHICAL SOCIETY, INDIA
(ESTD : 1962)



The Deccan Geographer



UGC Care Listed: A Peer Review and Refereed Journal

Vol. 61

No. 1 June and No. 2 December 2023

(ISSN-0011-7269)

CONTENTS

- | | | |
|-----|--|-----|
| (1) | Machine Learning and Quantum GIS for Flood Inundation Mapping of Gorakhpur District using Sentinel-1 and Sentinel-2 Radar Interferometry
–Dr. Swarnima Singh | 01 |
| (2) | Spatio-Temporal Pattern of Self-Reported Morbidity in Haryana
–Hemant and Dr. Rajeshwari | 15 |
| (3) | Vulnerability of Erosion in Rohmoria River Bank of Upper Brahmaputra Valley, Assam
–Lonkham Boruah | 29 |
| (4) | Assessment of Mgnrega in Nuh District of Haryana
–Deepak Moda and Dr. Mehtab Singh | 44 |
| (5) | Agricultural Productivity in Haryana during Post Liberalisation Period- An Agro-Ecoregion Level Study
–Bhim Singh and Dr. M.S. Jaglan | 56 |
| (6) | Variability in Cropped Area and Floriculture in District Kannauj, Uttar Pradesh
–Praveen Kumar | 71 |
| (7) | Assessment of Land Use/Land Cover and Associated Environmental Indices using GIS and Remote Sensing Techniques
–Sushil Kumar Yadav, Himanshu Shekher, Pankaj Kumar Yadav and Dr. K.P. Goswami | 81 |
| (8) | Spatial Pattern of Agricultural Productivity in Mirzapur District, Uttar Pradesh
–Ankit Pandey, Nitin Kumar Mishra and Suleman | 93 |
| (9) | Increasing Incidences of Acute Diarrhoeal Diseases in Sikkim, India
–Dilli Ram Dahal, Vimal Khawas and Sundeep Chettri | 105 |

Contd...

(10)	Temporal Pattern of Cold Wave in Eastern Uttar Pradesh – <i>Rachana Mishra and Dipak Prasad</i>	116
(11)	Solid Waste Management and People's Perceptions in Kalimpong Municipality, West Bengal, India – <i>Anindita Barman and Professor Ranjan Roy</i>	125
(12)	Awareness among Women about Health Schemes and Services related to NRHM in Rural Areas of Haryana – <i>Dr. Mukesh Kumar</i>	137
(13)	Extraction of Built-up Area and Land Surface Temperature Estimation for Gurugram Metro Area, Haryana, India – <i>Dr. Rashmi Rani Anand, Prof. Kavita Arora and Ms. Sangini Verma</i>	149
(14)	Spatial Structure and Pattern of Urban Growth in Mysore Local Planning District Using Landscape Metrics – <i>Manjunatha, C. S., Dhanaraj, K. and Chandrashekara, B.</i>	162
(15)	Geospatial Technology Based Orchard Mapping and Assessment in Morshi Taluka, Maharashtra State – <i>Dr. Jyoti S. Rokde</i>	172

The Deccan Geographical Society of India
Department of Geography
Savitribai Phule Pune University, Pune (Maharashtra)



AWARENESS AMONG WOMEN ABOUT HEALTH SCHEMES AND SERVICES RELATED TO NRHM IN RURAL AREAS OF HARYANA

Dr. Mukesh Kumar

Abstract

(Revised Received in October, 2023)

Indian government launched National Rural Health Mission (NRHM) programme on 12th April 2005 to provide comprehensive and effective primary health care to the unprivileged and vulnerable sections of the society particularly children and women by enhancing access, availability and quality of public health services. So, present study tried to find out about the awareness about different health schemes and services related to NRHM in rural Haryana. Study found that only 25 percent women heard about NRHM. Study found that about 90 percent surveyed mothers have heard about ASHA. But there is variation between village having health center and village having no health center. 92.1 percent mothers were aware about ASHA who were living in village which have a health center while 85 percent women were aware about ASHA who were residing in a village which have no health center. Study also found that only 36.4 percent women were aware about Janani Surakshya Yojana (JSY) scheme. But there is variation in this between women lives in a village which have health center (40 percent) and women lives in a village which have no health center (30 percent). So it is clear that awareness about JSY scheme is more in those women who are residing in a village which have a health center. Study found that there were only 26.8 percent women were registered under JSY scheme among the surveyed women. It is also noticeable that women residing in a village which have a health center (58.5 percent) are more than double registered than women residing in a village which have no health center (27.8 percent) in first trimester. So, it is found that the awareness about NRHM related scheme in the study area is very low; which shows that there is low impact of NRHM in the studied rural area of Haryana.

Introduction

Health is a fundamental human right and it is the responsibility of the Government's to provide health care to all people in equal proportion. Ever since India's independence in 1947, various national schemes and programmes have been

References

- Das, A. and Srivastva, H.C. (2009): "Maternal Health Care Utilization Among The Tribals of Madhya Pradesh: Constraints, Prospectus And Ramification for Well-Being", *Man In India*, Vol.89, No.3, pp. 239-250.
- Dasgupta, Rajib and Bisht, Ramila (2010): "The Missing Mission of Health", *Economic and Political Weekly*, Vol.XLV, No. 6, pp.16-18.
- Government of Haryana (2012) Statistical Abstract 2010-11, Economic and Statistical Adviser, Haryana.
- Gupta S.K, Pal D.K, Tiwari R., Garg R. , Sarawagi R., Shrivastava A.K. and Gupta P.(2011): "Assessment of Janani Suraksha Yojana (JSY) – in Jabalpur, Madhya Pradesh: knowledge, attitude and utilization pattern of beneficiaries: a descriptive study", *International Journal of Current Biological and Medical Science*, Vol.1, No.2, pp. 06-11.
- Husain, Zakir (2011): "Health of National Rural Health Mission", *Economic and Political Weekly*, Vol. XLVI, No. 4, pp. 53-60.
- Joshi, Sujay, R and George, Mathew (2012): "Healthcare through Community Participation: Role of ASHAs", *Economic and Political Weekly*, Vol. XLVII, No. 10, pp. 70-76.
- Mahadevia, Darshini (2000): "Health for All in Gujarat: Is It Achievable?", *Economic and Political Weekly*, Vol.35, No. 35/36, pp. 3193-3204.
- Malik J.S, Kalhan M., Punia A., Sachdeva S. and Behera B.K. (2013): "Utilization of Health Services under Janani Suraksha Yojna in Rural Haryana", *International Journal of Medicine and Public Health*, Vol. 3, No. 3, pp. 176-179.
- Mehrotra, S. (2008): "Public Health System in UP: What Can Be Done?", *Economic and Political Weekly*, Vol. 43 No. 49, pp. 46-53.
- Mousumi Datta and Nirmalya Manna (2012): "A study on socio-demographic correlates of maternal health care utilization in a rural area of West Bengal, India", *Global Journal of Medicine and Public Health*, Vol. 1, No. 4, pp. 7-12.
- National Rural Health Mission (2009): Four Years of NRHM, Ministry of Health and Family Welfare, Government of India, New Delhi.
- Tandon, B.B. and Mehra, Rajneesh (2011): "Towards Health For All: Some Suggestions To policy Makers", *Indian Journal of Public Administrator*, Vol. 57, No. 1, pp. 66-81.
- Vineeth, S. (2010): "NRHM: Enabling Health For All", *Kurukshetra*, Vol. 58, No. 4, pp. 7-8.

--Dr. Mukesh Kumar
Assistant Professor in Geography
G.C.W Badhra, Ch. Dadri (Haryana)



Structural, dielectric and magnetic characteristics of Mn-substituted $\text{Bi}_{0.80}\text{Nd}_{0.20}\text{FeO}_3$ multiferroics

Sandhaya Jangra¹ · Sujata Sanghi² · Ashish Agarwal² · Satish Khosa³ · Manisha Rangil⁴

Received: 17 March 2021 / Accepted: 9 June 2021

© The Author(s), under exclusive licence to Springer-Verlag GmbH, DE part of Springer Nature 2021

Abstract

Recently, investigations have been focused on modified BiFeO_3 multiferroics for obtaining improved physical properties. Keeping this in view, multiferroics having composition $\text{Bi}_{0.80}\text{Nd}_{0.20}\text{Fe}_{1-x}\text{Mn}_x\text{O}_3$ ($x=0.01, 0.03$ and 0.05) have been synthesized using solid-state reaction method. The crystal structure was examined by X-ray diffraction technique and Rietveld refinement. All unit cells were described by a combination of rhombohedral ($R3c$) and orthorhombic ($Pbnm$) phase. Increasing Mn content results in phase transformation with increasing orthorhombic character and major contribution by rhombohedral symmetry for all samples. Dielectric measurements carried out by impedance/gain phase analyzer in the frequency range from 10 Hz to 1 MHz. Dielectric constant (ϵ') and loss tangent ($\tan \delta$) show dispersion in the lower frequency range. The characteristics of the Nyquist plot confirmed the non-Debye type of relaxation processes with negative temperature coefficient behaviour of resistance (NTCR) in the ceramics. A strong variation in impedance is observed with Mn content also confirmed by conductivity analysis. The variation of ' s ' with temperature described that conduction mechanism is overlapping of large polaron tunnelling for $x=0.01, 0.03$ and for $x=0.05$ the appropriate mechanism is small polaron tunnelling. Magnetic measurements indicate the change in ferromagnetic character which might be due to a small change in structural parameters and maximum remanent magnetization is $M_r=0.023$ emu/g and coercive field is $H_c=0.760$ kOe for $x=0.05$. These prepared materials with improved multiferroic properties may lead to many technical applications, such as sensors, transducers and memory devices, etc.

Keywords Multiferroics · Perovskite · Phase transition · Rietveld refinement · Dielectric properties

1 Introduction

The best approaches to create novel materials with rich functional properties can be achieved by combining two or more physical properties, viz., electrical, magnetism and elasticity, in a single material [1]. Among various types of materials, multiferroics possess such characteristics. Multiferroics have a combination of ferromagnetic, ferroelectric

and ferroelastic orders in a single material and gained importance due to their fascinating properties [2, 3]. Along with multiferroic materials, nano-sized materials have been drawing attention in such potential applications [4–6]. These types of multifunctional properties can be attained by either naturally occurring multiple ferroic orders or preparing composite using two separate single ferroic order materials [7, 8]. These materials have an additional property known as magneto-electric coupling (M–E coupling) in which magnetization can be controlled by applying an electric field and vice versa [9, 10]. This magnetoelectric effect contributes to an additional degree of freedom that supports rich functional properties in multiferroic materials [4]. On the basis of origin of these ferroic orderings, multiferroic materials can be divided in two categories, such as: (1) Intrinsic multiferroics and (2) Artificial multiferroics. Intrinsic multiferroics are those natural materials that possess two ferroic orders in a single-phase material. The main drawbacks of such single-phase/intrinsic multiferroic materials are that they possess

✉ Sujata Sanghi
sujkash@yahoo.com; sanghi@gjust.org

¹ Government College for Women, Badhra (Charkhi Dadri), Haryana 127308, India

² Department of Physics, Guru Jambheshwar University of Science and Technology, Hisar, Haryana 125001, India

³ Materials Research Laboratory, Department of Physics, Deenbandhu Chhotu Ram University of Science and Technology, Murthal, Sonapat, Haryana 131039, India

⁴ Vaish College, Rohtak, Haryana 124001, India

REVIEW

A minireview of 1,2,3-triazole hybrids with O-heterocycles as leads in medicinal chemistry

Bhavna Saroha¹ | Gourav Kumar¹ | Ramesh Kumar¹ | Meena Kumari² | Suresh Kumar¹

¹Department of Chemistry, Kurukshetra University, Kurukshetra, India

²Department of Chemistry, Govt. College for Women Badhra, Charkhi Dadri, India

Correspondence

Suresh Kumar, Department of Chemistry, Kurukshetra University, Kurukshetra, India.
Email: suresh_dua47@rediffmail.com

Funding information

Council of Scientific and Industrial Research, India; UGC New Delhi; Haryana State Council for Science and Technology

Abstract

Over the past few decades, the dynamic progress in the synthesis and screening of heterocyclic compounds against various targets has made a significant contribution in the field of medicinal chemistry. Among the wide array of heterocyclic compounds, triazole moiety has attracted the attention of researchers owing to its vast therapeutic potential and easy preparation via copper and ruthenium-catalyzed azide-alkyne cycloaddition reactions. Triazole skeletons are found as major structural components in a different class of drugs possessing diverse pharmacological profiles including anti-cancer, anti-bacterial, anti-fungal, anti-viral, anti-oxidant, anti-inflammatory, anti-diabetic, anti-tubercular, and anti-depressant among various others. Furthermore, in the past few years, a significantly large number of triazole hybrids were synthesized with various heterocyclic moieties in order to gain the added advantage of the improved pharmacological profile, overcoming the multiple drug resistance and reduced toxicity from molecular hybridization. Among these synthesized triazole hybrids, many compounds are available commercially and used for treating different infections/disorders like tazobactam and cefatrizine as potent anti-bacterial agents while isavuconazole and ravuconazole as anti-fungal activities to name a few. In this review, we will summarize the biological activities of various 1,2,3-triazole hybrids with copious oxygen-containing heterocycles as lead compounds in medicinal chemistry. This review will be very helpful for researchers working in the field of molecular modeling, drug design and development, and medicinal chemistry.

KEYWORDS

1,2,3-triazole, biological activities, hybrid compounds, oxygen heterocycles, structure-activity relationship

1 | INTRODUCTION

In order to grapple with the diseases originating from various microbes (bacterial/viral/others), their mutated forms and resistant strains as well as from various bodily malfunctions and mutated genes in humans/other organisms, there remains the continuous need for

biologically potent chemical entities capable of treating these disorders. Heterocyclic compounds form the backbone of medicinal chemistry. The great breakthroughs in the synthesis of various complex heterocyclic skeletons have made possible, the task of creating the libraries of unique heterocyclic compounds. Furthermore, the screening of these huge numbers of heterocyclic

- as selective *Mycobacterium tuberculosis* dihydrofolate reductase inhibitors with potent whole cell activity. *European Journal of Medicinal Chemistry*, 144, 262–276. <https://doi.org/10.1016/j.ejmech.2017.12.017>
- Zhang, B. (2019). Comprehensive review on the anti-bacterial activity of 1, 2, 3-triazole hybrids. *European Journal of Medicinal Chemistry*, 168, 357–372. <https://doi.org/10.1016/j.ejmech.2019.02.055>
- Zhang, H.-Z., Zhao, Z.-L., & Zhou, C.-H. (2018). Recent advance in oxazole-based medicinal chemistry. *European Journal of Medicinal Chemistry*, 144, 444–492. <https://doi.org/10.1016/j.ejmech.2017.12.044>
- Zhang, S., Xu, Z., Gao, C., Ren, Q.-C., Chang, L., Lv, Z.-S., & Feng, L.-S. (2017). Triazole derivatives and their anti-tubercular activity. *European Journal of Medicinal Chemistry*, 138, 501–513. <https://doi.org/10.1016/j.ejmech.2017.06.051>
- Zhou, Y. X., Zhang, H., & Peng, C. (2014). Puerarin: a review of pharmacological effects. *Phytotherapy Research*, 28(7), 961–975. <https://doi.org/10.1002/ptr.5083>
- Zimmermann, L. A., de Moraes, M. H., da Rosa, R., de Melo, E. B., Paula, F. R., Schenkel, E. P., Steindel, M., & Bernardes, L. S. (2018). Synthesis and SAR of new isoxazole-triazole bis-heterocyclic compounds as analogues of natural lignans with antiparasitic activity. *Bioorganic & Medicinal Chemistry*, 26(17), 4850–4862. <https://doi.org/10.1016/j.bmc.2018.08.025>

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Saroha, B., Kumar, G., Kumar, R., Kumari, M., & Kumar, S. (2021). A minireview of 1;2,3-triazole hybrids with O-heterocycles as leads in medicinal chemistry. *Chemical Biology & Drug Design*, 00, 1–27. <https://doi.org/10.1111/cbdd.13966>

NEW Chemical Biology And Drug Design

Overview

Impact Factor

2.817



H Index

82

Google
scholar

Impact Factor

2.987

Scopus

I. Basic Journal Info

Country



United Kingdom (<https://www.scijournal.org/country/united-kingdom>)

Journal ISSN: 17470277

Publisher: Blackwell (<https://www.scijournal.org/publishers/blackwell>)

History: 2006-ongoing

Journal Homepage: Link (<http://www.wiley.com/bw/journal.asp?ref=1747-0277>)

How to Get Published:



Review

Recent Developments in Nanocatalyzed Green Synthetic Protocols of Biologically Potent Diverse O-Heterocycles—A Review

Suresh Kumar ^{1,*}, Bhavna Saroha ¹, Gourav Kumar ¹, Ekta Lathwal ^{1,2}, Sanjeev Kumar ¹, Badri Parshad ³, Meena Kumari ⁴, Naveen Kumar ⁵, Mabel M. Mphahlele-Makgwane ^{6,*} and Peter R. Makgwane ^{7,8}

- ¹ Department of Chemistry, Kurukshetra University, Kurukshetra 136119, Haryana, India; bhavnasaroha0712@gmail.com (B.S.); gouravbura1994@gmail.com (G.K.); ekta.lathwal@gmail.com (E.L.); sanjeevkumarsahrawat@gmail.com (S.K.)
 - ² Department of Chemistry, Govt. College, Tigaon, Faridabad 121101, Haryana, India
 - ³ Wellman Center for Photomedicine, Massachusetts General Hospital, Harvard Medical School, Charlestown, MA 02129, USA; bparshad@mgh.harvard.edu
 - ⁴ Department of Chemistry, Govt. College for Women, Badhra, Charkhi Dadri 127306, Haryana, India; jakhar7meena@gmail.com
 - ⁵ Department of Chemistry, Maharishi Dayanand University, Rohtak 124001, Haryana, India; naveenkumar.chem@mdurohtak.ac.in
 - ⁶ Department of Water and Sanitation, University of Limpopo, Private Bag X1106, Sovenga 0727, South Africa
 - ⁷ Centre for Nanostructures and Advanced Materials, Council for Scientific and Industrial Research (CSIR), Pretoria 0001, South Africa; pmakgwane@csir.co.za
 - ⁸ Department of Chemistry, University of the Western Cape, Bag X17, Robert Sobukwe Drive, Bellville 7535, South Africa
- * Correspondence: suresh_dua47@rediffmail.com (S.K.); mabel.mphahlele-makgwane@ul.ac.za (M.M.M.-M.)



Citation: Kumar, S.; Saroha, B.; Kumar, G.; Lathwal, E.; Kumar, S.; Parshad, B.; Kumari, M.; Kumar, N.; Mphahlele-Makgwane, M.M.; Makgwane, P.R. Recent Developments in Nanocatalyzed Green Synthetic Protocols of Biologically Potent Diverse O-Heterocycles—A Review. *Catalysts* **2022**, *12*, 657. <https://doi.org/10.3390/catal12060657>

Academic Editor: Antonella Glisenti

Received: 4 May 2022

Accepted: 24 May 2022

Published: 15 June 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: The dynamic growth in green organic synthetic methodologies for diverse heterocyclic scaffolds has substantially contributed to the field of medicinal chemistry over the last few decades. The use of hybrid metal nanocatalysts (NCs) is one such benign strategy for ensuring the advancement of modern synthetic chemistry by adhering to the principles of green chemistry, which call for a sustainable catalytic system that converts reacting species into profitable chemicals at a faster rate and tends to reduce waste generation. The metal nanoparticles (NPs) enhance the exposed surface area of the catalytic active sites, thereby making it easier for reactants and metal NCs to have an effective interaction. Several review articles have been published on the preparation of metal NCs and their uses for various catalytic heterocyclic transformations. This review will summarize different metal NCs for the efficient green synthesis of various O-heterocycles. Furthermore, the review will provide a concise overview of the role of metal NCs in the synthesis of O-heterocycles and will be extremely useful to researchers working on developing novel green and simple synthetic pathways to various O-heterocyclic-derived molecules.

Keywords: nanocatalyst; green chemistry; chalcone; coumarin; furan; oxazole; pyran

1. Introduction

Chemistry is an undeniably important part of our daily lives, and its advancements have been developed to provide people with access to innovative medicines for healthier living conditions. Several heterocycle scaffolds have been found to be pharmaceutically active for applications in the development and formulation of new drugs [1]. Heterocycles are found in both natural and synthesized biologically active compounds [2]. Despite incidents [3] such as the clioquinol tragedy, the thalidomide tragedy, and others, the importance of heterocycles in drug discovery continues to rise due to their selectively pharmacophore binding potential. The heterocyclic skeleton can be found in a large number of commercially accessible Food and Drug Administration (FDA)-approved synthesized

90. Fekri, L.Z.; Nikpassand, M.; Shariati, S.; Aghazadeh, B.; Zarkeshvari, R.; Norouz pour, N. Synthesis and characterization of amino glucose-functionalized silica-coated NiFe_2O_4 nanoparticles: A heterogeneous, new and magnetically separable catalyst for the solvent-free synthesis of 2,4,5-trisubstituted imidazoles, benzo[d]imidazoles, benzo[d]oxazoles and azo-linked benzo[d]oxazoles. *J. Organomet. Chem.* **2018**, *871*, 60–73.
91. Aboonajmi, J.; Sharghi, H.; Aberi, M.; Shiri, P. Consecutive Oxidation/Condensation/Cyclization/Aromatization Sequences Catalyzed by Nanostructured Iron(III)-Porphyrin Complex towards Benzoxazole Derivatives. *Eur. J. Org. Chem.* **2020**, *2020*, 5978–5984. [CrossRef]
92. Yu, C.; Guo, X.; Shen, B.; Xi, Z.; Li, Q.; Yin, Z.; Liu, H.; Muzzio, M.; Shen, M.; Li, J.; et al. One-pot formic acid dehydrogenation and synthesis of benzene-fused heterocycles over reusable AgPd/WO_2 nanocatalyst. *J. Mater. Chem. A* **2018**, *6*, 23766–23772. [CrossRef]
93. Shanshak, M.; Budagumpi, S.; Malecki, J.G.; Keri, R.S. Green synthesis of 3, 4-disubstituted isoxazol-5 (4H)-ones using $\text{ZnO}/\text{Fe}_3\text{O}_4$ core-shell nanocatalyst in water. *Appl. Organomet. Chem.* **2020**, *34*, e5544. [CrossRef]
94. SUTHAR, M.; KUMBHANI, J.; BHATT, K.D. Pyran Heterocyclic Compound as the Prosperous Scaffolds for Biological Sensor (A-Review). *Orient. J. Chem.* **2021**, *37*, 1280–1286. [CrossRef]
95. Safari, F.; Hosseini, H.; Bayat, M.; Ranjbar, A. Synthesis and evaluation of antimicrobial activity, cytotoxic and pro-apoptotic effects of novel spiro-4 H-pyran derivatives. *RSC Adv.* **2019**, *9*, 24843–24851. [CrossRef]
96. Yang, Z.-J.; Gong, Q.-T.; Wang, Y.; Yu, Y.; Liu, Y.-H.; Wang, N.; Yu, X.-Q. Biocatalytic tandem multicomponent reactions for one-pot synthesis of 2-Amino-4H-Pyran library and in vitro biological evaluation. *Mol. Catal.* **2020**, *491*, 110983. [CrossRef]
97. Sirous, H.; Chemi, G.; Gemma, S.; Butini, S.; Debyser, Z.; Christ, F.; Saghale, L.; Brogi, S.; Fassihi, A.; Campiani, G.; et al. Identification of Novel 3-Hydroxy-pyran-4-One Derivatives as Potent HIV-1 Integrase Inhibitors Using in silico Structure-Based Combinatorial Library Design Approach. *Front. Chem.* **2019**, *7*, 574. [CrossRef] [PubMed]
98. Arustamyan, Z.S.; Margaryan, R.E.; Aghekyan, A.A.; Panosyan, G.A.; Muradyan, R.E.; Tumajyan, A.E. Synthesis and Anti-Inflammatory Properties of Substituted 5-(Tetrahydro-4-phenyl-2H-pyran-4-yl)-4H-1,2,4-triazole-3-thiols. *Russ. J. Org. Chem.* **2021**, *57*, 195–202. [CrossRef]
99. Bhat, Z.S.; Lah, H.U.; Rather, M.A.; Maqbool, M.; Ara, T.; Ahmad, Z.; Yousuf, S.K. Synthesis and in vitro evaluation of substituted 3-cinnamoyl-4-hydroxy-pyran-2-one (CHP) in pursuit of new potential antituberculosis agents. *MedChemComm* **2018**, *9*, 165–172. [CrossRef]
100. Sajjadi-Ghotbabadi, H.; Javanshir, S.; Rostami-Charati, F.; Zahra Sayyed-Alangi, S. Eco-compatible synthesis of novel 3-hydroxyflavones catalyzed by KF-impregnated mesoporous natural zeolite clinoptilolite. *Chem. Heterocycl. Compd.* **2018**, *54*, 508–513. [CrossRef]
101. Ghasemzadeh, M.S.; Akhlaghinia, B. $\gamma\text{-Fe}_2\text{O}_3/\text{SiO}_2\text{-EC-ZnII}$: A Magnetic Recyclable Nanocatalyst for the Synthesis of Spiro [indoline-3, 9'-xanthene] trione Derivatives in Aqueous Media. *ChemistrySelect* **2018**, *3*, 3161–3170. [CrossRef]
102. Mohammadi, H.; Shaterian, H.R. $\gamma\text{-Fe}_2\text{O}_3/\text{SiO}_2\text{-}\gamma\text{-aminobutyric acid}$ as a novel superparamagnetic nanocatalyst promoted green synthesis of chromeno[4,3,2-de][1,6]naphthyridine derivatives. *Monats. Chem.* **2019**, *150*, 327–337. [CrossRef]
103. Eshtehardian, B.; Rouhani, M.; Mirjafary, Z. Green protocol for synthesis of MgFe_2O_4 nanoparticles and study of their activity as an efficient catalyst for the synthesis of chromene and pyran derivatives under ultrasound irradiation. *J. Iran. Chem. Soc.* **2020**, *17*, 469–481. [CrossRef]
104. Pourshojaei, Y.; Zolala, F.; Eskandari, K.; Talebi, M.; Morsali, L.; Amiri, M.; Khodadadi, A.; Shamsimeymandi, R.; Faghih-Mirzaei, E.; Asadipour, A. Nickel ferrite (NiFe_2O_4) nanoparticles as magnetically recyclable nanocatalyst for highly efficient synthesis of 4H-chromene derivatives. *J. Nanosci. Nanotechnol.* **2020**, *20*, 3206–3216. [CrossRef] [PubMed]
105. Singh, P.; Yadav, P.; Mishra, A.; Awasthi, S.K. Green and Mechanochemical One-Pot Multicomponent Synthesis of Bioactive 2-amino-4H-benzo[b]pyrans via Highly Efficient Amine-Functionalized $\text{SiO}_2/\text{Fe}_3\text{O}_4$ Nanoparticles. *ACS Omega* **2020**, *5*, 4223–4232. [CrossRef] [PubMed]
106. Lati, M.P.; Shirini, F.; Alinia-Asli, M.; Rezvani, M.A. $\text{Fe}_3\text{O}_4/\text{SiO}_2\text{-Sultone}$: A Novel and Recyclable Magnetic Nanocatalyst for the Efficient Synthesis of 3, 4 Dihydropyrano [c] Chromenes and 2-Amino-4H-Chromene Derivatives. *J. Nanosci. Nanotechnol.* **2020**, *20*, 973–982. [CrossRef]
107. Karami, S.; Dekamin, M.G.; Valiey, E.; Shakib, P. DABA MNPs: A new and efficient magnetic bifunctional nanocatalyst for the green synthesis of biologically active pyrano[2,3-c]pyrazole and benzylpyrazolyl coumarin derivatives. *New J. Chem.* **2020**, *44*, 13952–13961. [CrossRef]
108. Kamalzare, M.; Ahghari, M.R.; Bayat, M.; Maleki, A. $\text{Fe}_3\text{O}_4/\text{chitosan-tannic acid}$ bionanocomposite as a novel nanocatalyst for the synthesis of pyranopyrazoles. *Sci. Rep.* **2021**, *11*, 20021. [CrossRef]
109. Hoseinzade, K.; Mousavi-Mashhadi, S.A.; Shiri, A. An efficient and green one-pot synthesis of tetrahydrobenzo[a]xanthenes, 1,8-dioxo-octahydroxanthenes and dibenzo[a,j]xanthenes by $\text{Fe}_3\text{O}_4/\text{Agar-Ag}$ as nanocatalyst. *Mol. Divers.* **2022**. [CrossRef]
110. Tanuraghaj, H.M.; Farahi, M. A novel protocol for the synthesis of pyrano[2,3-h]coumarins in the presence of $\text{Fe}_3\text{O}_4/\text{SiO}_2/(\text{CH}_2)_3\text{-OCO}_2\text{Na}$ as a magnetically heterogeneous catalyst. *New J. Chem.* **2019**, *43*, 4823–4829. [CrossRef]

Catalysts Impact Factor & Key Scientometrics

NEW Catalysts Overview

Impact Factor

H Index

Impact Factor

<https://www.scijournal.org/impact-factor-of-catalysts.shtml>

1/15

3/25/24, 12:47 AM

Catalysts - SCI Journal



Web of
Science
Group



Google



Scopus

I. Basic Journal Info

Country



Switzerland (<https://www.scijournal.org/country/switzerland>)

Journal ISSN: 20734344

Publisher: Multidisciplinary Digital Publishing Institute (MDPI)
(<https://www.scijournal.org/publishers/multidisciplinary-digital-publishing-institute-mdpi>)

History: 2011-ongoing

Journal Homepage: Link (NA)

How to Get Published:

Find out more (/articles)

* Research Categories

Design, synthesis, biological evaluation, and molecular docking studies of some novel *N,N*-dimethylaminopropoxy-substituted aurones

Gourav Kumar¹ | Bhavna Saroha¹ | Ramesh Kumar¹ | Meena Kumari² | Sunita Dalal³ | Suresh Kumar¹

¹Department of Chemistry, Kurukshetra University, Kurukshetra, India

²Department of Chemistry, Govt. College for Women, Badhra, Charkhi Dadri, India

³Department of Biotechnology, Kurukshetra University, Kurukshetra, India

Correspondence

Suresh Kumar, Department of Chemistry, Kurukshetra University, Kurukshetra, Haryana, 136119, India.
Email: suresh_dua47@rediffmail.com

Funding information

Council of Scientific and Industrial Research, India; Haryana State Council for Science and Technology, Grant/Award Number: HSCST/R&D/2018/2097; University Grants Commission

Abstract

In continuation of our ongoing research on the discovery of novel and potentially bioactive aurones, we have designed and synthesized some novel *N,N*-dimethylaminopropoxy-substituted pyrazole-based aurones **10(a-l)**. These pyrazole-benzofuranone hybrid compounds were characterized by using their IR, ¹H-NMR, ¹³C-NMR, and mass spectrometry data. Compound **10c** was used as a model to further explicate the structure of tilted compounds by means of ¹H-¹H COSY, ¹H-¹³C HMQC, ¹H-¹³C HMBC, ¹H-¹H TOCSY, ¹H-¹H NOSEY, DEPT-45°, DEPT-90°, and DEPT-135° NMR spectra. The comparative molecular docking study of *N,N*-dimethylaminopropoxy-substituted pyrazole-based aurones and standard drugs (Ampicillin and Chloramphenicol) against *Bacillus subtilis* (PDB: 6tzip) active site was performed to determine the binding interactions, binding energy, and orientation of the molecules at the active site of the target protein. Out of these synthesized compounds, five best analogs (**10b**, **10f**, **10h**, **10k**, and **10l**) of docking results were also evaluated for their in vitro antibacterial potential against *Bacillus subtilis* to validate the docking results.

1 | INTRODUCTION

Aurone, that is, 2-benzylidenebenzofuran-3(2*H*)-one, is a ubiquitous family of compounds having a benzofuranone heterocyclic ring connected to phenyl moiety via a carbon-carbon exocyclic double bond. These fascinating scaffolds are secondary plant metabolites and are liable for imparting pigmentation in flowers, fruits, and other colored portions of the plants [1]. Aurones are the active ingredients of various conventional remedies like *Vaccinium oxycoccos* (European cranberry), *Glycyrrhiza glabra* (Licorice), and *Ceanothus americanus* (New Jersey tea) [2]. Even though aurones have very confined existence in nature, they constitute a blazing class of pharmacologically effectual skeletons, which exhibit prodigious

range of biological accent [3] including anticancer [4], antioxidant [5], antiparasitic [6], antiobesity [7], tyrosinase inhibition [8], cathepsin B inhibition [9], anti-hormonal [7], antileishmanial [10], antiquorum sensing [11], anti-Alzheimer [12], and antiinflammatory [13] activities. Some of the eminent aurones are cephalocerone, hemiltron, hispidol, and maritimetin, where Cephalocerone, extracted from the plant of *Cephalocereus senilis*, showed antibacterial activity [14]; hamiltrone, isolated from plant *Uvaria hamiltonii* displays DNA strand scission activity [15]; hispidol obtained from soybean *Glycine max* exhibits antifungal activity [16]; and maritimetin present in genus *Coreopsis* depicted anticarcinogen and antioxidant activities [7,17]. Figure 1.

- Lett.* **2004**, *14*, 4307. (c) Y. H. Lee, M. C. Shin, Y. D. Yun, S. Y. Shin, J. M. Kim, J. M. Seo, N.-J. Kim, J. H. Ryu, Y. S. Lee, *Bioorg. Med. Chem.* **2015**, *23*, 231. (d) C.-H. Tseng, Y.-L. Chen, K.-Y. Chung, C.-M. Cheng, C.-H. Wang, C.-C. Tzeng, *Bioorg. Med. Chem.* **2009**, *17*, 7465.
- [34] G. Kumar, E. Lathwal, B. Saroha, S. Kumar, S. Kumar, N. S. Chauhan, T. Kumar, *ChemistrySelect* **2020**, *5*, 3539.
- [35] S. Kumar, E. Lathwal, G. Kumar, B. Saroha, S. Kumar, S. Mahata, P. K. Sahoo, V. D. Nasare, *Chem. Data Collect.* **2020**, *30*, 100559.
- [36] (a) H. B'Bhatt, S. Sharma, J. Arabian, *Chem* **2017**, *10*, S1590. (b) P. A. Elzahhar, R. A. Alaaeddine, R. Nassra, A. Ismail, H. F. Labib, M. G. Temraz, A. S. F. Belal, A. F. El-Yazbi, *J. Enzyme Inhib. Med. Chem.* **2021**, *36*, 669. (c) O. Prakash, R. Kumar, V. Parkash, *Eur. J. Med. Chem.* **2008**, *43*, 435. (d) O. Prakash, K. Pannu, R. Naithani, H. Kaur, *Synth. Commun.* **2006**, *36*, 3479. (e) O. Prakash, D. K. Aneja, S. Arora, C. Sharma, K. R. Aneja, *Med. Chem. Res.* **2012**, *21*, 10. (f) E. V. Rudyakova, V. A. Savosik, L. K. Papernaya, A. I. Albanov, I. T. Evstafeva, G. G. Levkovskaya, *Russ. J. Org. Chem.* **2009**, *45*, 1040. (g) N. Harikrishna, A. M. Isloor, R. Ananda, A. Obaid, H.-K. Fun, *RSC Adv.* **2015**, *5*, 43648–43659.
- [37] X. Zheng, H. Wang, Y. M. Liu, X. Yao, M. Tong, Y. H. Wang, D. F. Liao, *J. Heterocyclic Chem.* **2015**, *52*, 296.
- [38] S. Bai, P. Bharti, L. Seasotiya, A. Malik, S. Dalal, *Pharm. Biol.* **2015**, *53*, 326.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: G. Kumar, B. Saroha, R. Kumar, M. Kumari, S. Dalal, S. Kumar, *J. Heterocycl. Chem.* **2021**, *1*. <https://doi.org/10.1002/jhet.4384>

Impact Factor

2.193




H Index

NA



Impact Factor

NA



I. Basic Journal Info

Journal ISSN: NA

Publisher: NA (<https://www.scijournal.org/publishers/na>)

History: NA

Journal Homepage: Link (NA)


How to Get Published:

Find out more (/articles)

Research Categories

Scope/Description:

NA



Data Article

Synthesis of propynyloxy substituted some novel aurones as potent cathepsin B inhibitors



Bhavna Saroha^a, Gourav Kumar^a, Ekta Lathwal^{a,b}, Sanjeev Kumar^a,
Meena Kumari^c, Nitika Mor^a, Neera Raghav^a, Suresh Kumar^{a,*}

^a Department of Chemistry, Kurukshetra University, Kurukshetra, 136119, Haryana, India

^b T.D.L. Govt. College for Women, Murthal, Sonapat 131027, Haryana, India.

^c Govt. College for Women, Badhra, Charkhi Dadri, 127308, Haryana, India

ARTICLE INFO

Article history:

Received 20 October 2020

Revised 13 December 2020

Accepted 13 December 2020

Available online 18 December 2020

Keywords:

Aurone

Propargyl bromide

Synthesis

Cathepsin B inhibitors

Molecular docking studies

ABSTRACT

A two-step synthesis was designed to develop a novel series of 6-propynyloxyaurones, **2(a-r)**. The synthesized aurone derivatives were characterized by their IR, NMR, and mass spectrometry data. The present study proposes these aurones as anti-inflammatory and anti-Alzheimer candidates on the basis of their cathepsin B inhibitory activities. Anti-cathepsin B activity of some of the synthesized aurones was found, either equivalent to or higher than the reference drugs, aspirin and curcumin at submicromolar concentrations. Among the studied compounds, **2e**, **2i**, **2n**, **2q**, and **2r** exhibited much higher activity in comparison to the standard drugs. The molecular docking study also highlighted compound **2r** for its greater anti-cathepsin B activity among all the designed aurones.

© 2020 Elsevier B.V. All rights reserved.

Specifications Table

Subject area	Organic Chemistry
Compounds	Aurones
Data category	Spectral, Synthesis, Molecular docking
Data acquisition format	NMR, IR, Mass spectra
Data type	Analysis
Procedure	Brief description of the procedure (e.g., transformation, simulation, etc.)
Data accessibility	A series of novel aurones were synthesized and characterized by spectral data and further evaluated for their biological activity.

1. Rationale

Cathepsin B, the most abundantly expressed mammalian lysosomal protease, involved in intracellular protein catabolism, has emerged out as a potential drug target for a number of diseases [1]. The elevated level of cathepsin B has been found responsible for several pathological conditions, which include inflammation [2,3], cancer [4–6], neurological disorders such as Alzheimer's [3,7] and cardiac disorders [8] among various others. Elevated levels of cathepsins have been found vis-à-vis decreased levels of intracellular inhibitors [9] thereby generating the need for development of novel cathepsin B inhibitors.

* Corresponding author.

E-mail address: suresh_dua47@rediffmail.com (S. Kumar).

- [30] W.J. Geldenhuys, M.O. Funk, C.J. Van der Schyf, R.T. Carroll, A scaffold hopping approach to identify novel monoamine oxidase B inhibitors, *Bioorg. Med. Chem. Lett.* 22 (2012) 1380–1383, doi:10.1016/j.bmcl.2011.12.056.
- [31] S. Shrestha, S. Natarajan, J.-H. Park, D.-Y. Lee, J.-G. Cho, G.-S. Kim, Y.-J. Jeon, S.-W. Yeon, D.-C. Yang, N.-I. Baek, Potential neuroprotective flavonoid-based inhibitors of CDK5/p25 from *Rhus parviflora*, *Bioorg. Med. Chem. Lett.* 23 (2013) 5150–5154, doi:10.1016/j.bmcl.2013.07.020.
- [32] V. Zwick, A.-O. Chatzivasileiou, N. Deschamps, M. Roussaki, C.A. Simões-Pires, A. Nurisso, I. Denis, C. Blanquart, N. Martinet, P.-A. Carrupt, Aurones as histone deacetylase inhibitors: identification of key features, *Bioorg. Med. Chem. Lett.* 24 (2014) 5497–5501, doi:10.1016/j.bmcl.2014.10.019.
- [33] R. Shamsimeymani, Y. Pourshojaei, K. Eskandari, M. Mohammadi-Khanaposhtani, A. Abiri, A. Khodadadi, A. Langarizadeh, F. Sharififar, B. Amirheidari, T. Akbarzadeh, Design, synthesis, biological evaluation, and molecular dynamics of novel cholinesterase inhibitors as anti-Alzheimer's agents, *Arch. Pharm.* 352 (2019) 1800352, doi:10.1002/ardp.201800352.
- [34] X. Dong, J. Chen, C. Jiang, T. Liu, Y. Hu, Design, synthesis, and biological evaluation of prenylated chalcones as vasorelaxant agents, *Arch. Pharm.* 342 (2009) 428–432, doi:10.1002/ardp.200800229.
- [35] F. Jin, N. Zhang, C. Tan, D. Gao, C. Zhang, F. Liu, Z. Chen, C. Gao, H. Liu, S. Li, 2'-Chloro-4'-aminoflavone derivatives selectively targeting hepatocarcinoma cells: convenient synthetic process, G2/M cell cycle arrest and apoptosis triggers, *Arch. Pharm.* 345 (2012) 525–534, doi:10.1002/ardp.201100383.
- [36] K.M. Amin, A.M. Taha, R.F. George, N.M. Mohamed, F.F. Elsenduny, Synthesis, antitumor activity evaluation, and DNA-binding study of coumarin-based agents, *Arch. Pharm.* 351 (2018) 1700199, doi:10.1002/ardp.201700199.
- [37] N. Raghav, R. Kaur, Chalcones, semicarbazones and pyrazolines as inhibitors of cathepsins B, H and L, *Int. J. Biol. Macromol.* 80 (2015) 710–724, doi:10.1016/j.ijbiomac.2015.07.029.
- [38] I. Ravish, N. Raghav, SAR studies of differently functionalized 4'-phenylchalcone based compounds as inhibitors of cathepsins B, H and L, *RSC Adv* 5 (2015) 50440–50453, doi:10.1039/C5RA00357A.
- [39] L.M. de Novais, C.C. de Arceira, L.F. Ferreira, T.A. Ribeiro, P.T. Sousa Jr, M.J. Jacinto, M.G. de Carvalho, W.A. Judice, L.O. Jesus, A.A. de Souza, 4'-Hydroxy-6, 7-methylenedioxy-3-methoxyflavone: a novel flavonoid from *Dulacia eglei* with potential inhibitory activity against cathepsins B and L, *Fitoterapia* 132 (2019) 26–29, doi:10.1016/j.fitote.2018.08.005.
- [40] S.D. Ramalho, L.R. de Sousa, M.C. Burger, M.I.S. Lima, M.F.d.G.da Silva, J.B. Fernandes, P.C. Vieira, Evaluation of flavonols and derivatives as human cathepsin B inhibitor, *Nat. Prod. Res.* 29 (2015) 2212–2214, doi:10.1080/14786419.2014.1002404.
- [41] O.M. Bautista-Aguilera, A. Samadi, M. Chioua, K. Nikolic, S. Filipic, D. Agbaba, E. Soriano, L. de Andres, M.I. Rodríguez-Franco, S. Alcaro, N-methyl-N-((1-methyl-5-(3-(1-(2-methylbenzyl) piperidin-4-yl) propoxy)-1 H-indol-2-yl) methyl) prop-2-yn-1-amine, a new cholinesterase and monoamine oxidase dual inhibitor, *J. Med. Chem.* 57 (2014) 10455–10463, doi:10.1021/jm501501a.
- [42] F.T. Zindo, J. Joubert, S.F. Malan, Propargylamine as functional moiety in the design of multifunctional drugs for neurodegenerative disorders: MAO inhibition and beyond, *Future Med. Chem.* 7 (2015) 609–629, doi:10.4155/fmc.15.12.
- [43] B. Kumar, M. Kumar, A.R. Dwivedi, V. Kumar, Synthesis, Biological Evaluation and Molecular Modeling Studies of Propargyl-Containing 2, 4, 6-trisubstituted pyrimidine derivatives as potential anti-parkinson agents, *ChemMedChem* 13 (2018) 705–712, doi:10.1002/cmdc.201700589.
- [44] B. Kumar, A.R. Dwivedi, B. Sarkar, S.K. Gupta, S. Krishnamurthy, A.K. Mantha, J. Parkash, V. Kumar, 4, 6-Diphenylpyrimidine derivatives as dual inhibitors of monoamine oxidase and acetylcholinesterase for the treatment of Alzheimer's disease, *ACS Chem. Neurosci.* 10 (2018) 252–265, doi:10.1021/acschemneuro.8b00220.
- [45] B. Kumar, V. Kumar, V. Prashar, S. Saini, A.R. Dwivedi, B. Bajaj, D. Mehta, J. Parkash, V. Kumar, Dipropargyl substituted diphenylpyrimidines as dual inhibitors of monoamine oxidase and acetylcholinesterase, *Eur. J. Med. Chem.* 177 (2019) 221–234, doi:10.1016/j.ejmech.2019.05.039.
- [46] Y.A.R. Núñez, M. Norambuena, A.R.R. Bohorquez, A. Morales-Bayuelo, M. Gutiérrez, Efficient synthesis and antioxidant activity of novel N-propargyl tetrahydroquinoline derivatives through the cationic Povarov reaction, *Heliyon* 5 (2019) e02174, doi:10.1016/j.heliyon.2019.e02174.
- [47] X. Zheng, H. Wang, Y.M. Liu, X. Yao, M. Tong, Y.H. Wang, D.F. Liao, Synthesis, characterization, and anticancer effect of trifluoromethylated aurone derivatives, *J. Heterocycl. Chem.* 52 (2015) 296–301, doi:10.1002/jhet.1969.
- [48] S. Kumar, An improved one-pot and eco-friendly synthesis of aurones under solvent-free conditions, *Green Chem. Lett. Rev.* 7 (2014) 95–99, doi:10.1080/17518253.2014.895867.
- [49] N. Raghav, M. Singh, S. Garg, R. Kaur, S. Jangra, I. Ravish, Ion exchangers: a useful tool for separation and simultaneous purification of lysosomal cysteine proteinases, cathepsins B, H and L, *Int. J. Pharm. Sci. Res.* 6 (2015) 2944–2949, doi:10.13040/IJPSR.0975-8232.6(7).2944-49.
- [50] N. Raghav, M. Singh, SAR studies of some acetophenone phenylhydrazone based pyrazole derivatives as anticathepsin agents, *Bioorg. Chem.* 75 (2017) 38–49, doi:10.1016/j.bioorg.2017.08.006.
- [51] C. Ferroni, A. Pepe, Y.S. Kim, S. Lee, A. Guerrini, M.D. Parenti, A. Tesei, A. Zamagni, M. Cortesi, N. Zaffaroni, 1, 4-Substituted triazoles as nonsteroidal anti-androgens for prostate cancer treatment, *J. Med. Chem.* 60 (2017) 3082–3093, doi:10.1021/acs.jmedchem.7b00105.

Chemical Data Collections Impact Factor & Key Scientometrics

NEW Chemical Data Collections Overview

Impact Factor

H Index

Impact Factor

<https://www.scijournal.org/impact-factor-of-chemical-data-collections.shtml>

1/16

3/25/24, 12:53 AM

Chemical Data Collections - SCI Journal



I. Basic Journal Info

Country



Netherlands (<https://www.scijournal.org/country/netherlands>)

Journal ISSN: 24058300

Publisher: Elsevier BV (<https://www.scijournal.org/publishers/elsevier-bv>)

History: 2016-2021

Journal Homepage: Link (<https://www.journals.elsevier.com/chemical-data-collections>)

How to Get Published:

Find out more (/articles)

Research Categories

■ Organic & Supramolecular Chemistry

Recent Advances in Synthesis and Biological Assessment of Quinoline-Oxygen Heterocycle Hybrids

Gourav Kumar,^[a] Bhavna Saroha,^[a] Ramesh Kumar,^[a] Meena Kumari,^[b] and Suresh Kumar^{*[a]}

The exploration of hybrid molecules for targeting various diseases represents a highly promising approach which has showcased significant momentum worldwide, in the last two decades. Diversely substituted quinoline scaffolds present in naturally occurring bioactive compounds, signify their role in various biological processes and stimulated the research developments around it, for diverse therapeutic applications. With the ongoing search for suitable pharmacophores, the quinoline derivatives have emerged as versatile compounds possessing significant and broad spectrum of biological activities. Further, conjugation of quinoline with other bioactive

heterocycles containing oxygen yielded hybrid organic frameworks with improved therapeutic profile leading to an emerging paradigm and tremendous growth in the contemporary drug discovery regime. The synthesis of these hybrid skeletons with high efficiency still remains the most strenuous task and a lot of research work is being performed in this direction. In the present article, we have reviewed the synthetic protocols leading to quinoline-oxygen heterocycle hybrids and also highlighted the systematic and biological assessment of developed pharmacophores.

1. Introduction

In order to combat the increasing number of viral and microbial infections and other diseases, there is an intensive demand for developing the innovative formulations with suitable pharmacophores which could be synthesized using novel and highly efficient synthetic strategies. The requirement for newer therapeutic moieties further amplifies with the increasing cases of side-effects and drug resistance. The chemotherapeutic response of these heterocyclic moieties is generated by their affinity to serve as ligands towards various biological receptors. The therapeutic response of these heterocyclic moieties is significantly influenced by the presence of different substituents through the changes in electronic structure and binding affinity. Quinoline i.e. benzo[b]pyridine or 1-azaphthalene is an ubiquitous heterocyclic moiety found in numerous biologically active natural products serving as anti-malarial,^[1] antiparasitic,^[2] anti-cancer,^[3] anti-inflammatory,^[4] anti-tubercular,^[5] anti-biotic,^[6] tyrosinase PDGF-RTK,^[7] anti-fungal,^[8] analgesic,^[9] cardiovascular^[10] activities among various others. Some of the well known drugs having quinoline skeleton involve: quinine, chloroquine, primaquine, amodiaquine etc. as antimalarial; moxifloxacin, ciprofloxacin as anti-tubercular agents,^[11] pyriminyl as anthelmintic agent^[12]; and cinchocaine as an anesthetic,^[13] and various others (Figure 1). Owing to the high medicinal value and good solubility in most of the organic solvents as well as partial solubility in water, many research

groups are elaborating the research work related to synthesis and bioactivity evaluation of quinoline based skeletons with an aim to exploit their inherent properties.^[14]

Further, in order to gain the benefits from molecular hybridization, a modern strategy for drug design in medicinal chemistry, the conjugation of quinoline scaffolds with other oxygen based bioactive heterocyclic compounds has gained much attention in the last few years. The molecular hybridization involves the chemical conjugation fusion of two or more pharmacophores with specific structural domains into a hybrid skeleton with significant enhancement expectation of improved therapeutic profile. These hybrid molecules possessing diverse skeletons not only provide increased efficacy and safety but also help in overcoming the resistance on account of multiple modes of action as well as binding with multiple targets.^[15] Among the various oxygen containing heterocycles, furans and pyrans have gathered enormous attention as versatile and fruitful organic compounds with impressive medicinal history. These vital oxygen heterocyclic compounds have reflected interesting pharmacological behaviors such as anti-hyperglycemic,^[16] analgesic,^[17] anti-tumor and kinase Inhibitor,^[18] anti-microbial,^[19] anti-parasitic,^[20] anti-viral,^[21] anti-cancer^[22] and anti-inflammation^[23] activities among many others.

The naturally occurring hybrid skeletons having quinoline and oxygen based heterocyclic motifs have attracted prodigious attention from chemists and pharmacologists working in the field of natural products owing to their simple structures and noteworthy bioactivities. Some of such hybrid molecules obtained from microorganisms and flowering plants are shown in Figure 2 and found to possess several bioactivities. Graveoline, a quinoline based hybrid alkaloid extracted from *Vepris punctate* was found to demonstrate significant herbicidal activity and cytotoxicity against A2780 cell lines.^[24] Also, 2-(3,4-

[a] G. Kumar, B. Saroha, Dr. R. Kumar, Dr. S. Kumar
Department of Chemistry, Kurukshetra University, Kurukshetra, Haryana, India-136119
E-mail: suresh_dua47@rediffmail.com

[b] Dr. M. Kumari
Department of Chemistry, Govt. College for Women, Badhra, Charkhi Dadri, Haryana, India- 127308 *

- [54] C. L. Yang, C. H. Tseng, Y. L. Chen, C. M. Lu, C. L. Kao, M. H. Wu, C. C. Tzeng, *Eur. J. Med. Chem.* **2010**, *45*, 602–607.
- [55] C. Shi, J. Wang, H. Chen, D. Shi, *J. Comb. Chem.* **2010**, *12*, 430–434.
- [56] K. C. Majumdar, S. Ponra, D. Ghosh, A. Taher, *Synlett* **2011**, *2011*, 104–110.
- [57] A. T. Khan, D. K. Das, *Tetrahedron Lett.* **2012**, *53*, 2345–2351.
- [58] Y. Thigulla, T. U. Kumar, P. Trivedi, B. Ghosh, A. Bhattacharya, *ChemistrySelect* **2017**, *2*, 2718–2721.
- [59] Y. Lin, D. Xing, W. B. Wu, G. Y. Xu, L. F. Yu, J. Tang, Y. B. Zhou, J. Li, F. Yang, *Molecules* **2020**, *25*.
- [60] J. P. S. Ferreira, S. M. Cardoso, F. A. Almeida Paz, A. M. S. Silva, V. L. M. Silva, *New J. Chem.* **2020**, *44*, 6501–6509.

Submitted: March 11, 2021

Accepted: May 12, 2021

NEW ChemistrySelect

Overview

Impact Factor

2.109



H Index

41



Impact Factor

2.24



I. Basic Journal Info

Country



United Kingdom (<https://www.scijournal.org/country/united-kingdom>)

Journal ISSN: 23656549

Publisher: Wiley-Blackwell Publishing Ltd (<https://www.scijournal.org/publishers/wiley-blackwell-publishing-ltd>)

History: 2016-2021

Journal Homepage: Link (<https://onlinelibrary.wiley.com/journal/23656549>)

How to Get Published:



Ultrasound assisted a one pot multicomponent and greener synthesis of 1,2,3-triazole incorporated aurone hybrids: Cathepsin B inhibition, anti-cancer activity against AGS cell line, and *in-silico* docking evaluation

Bhavna Saroha^a, Gourav Kumar^a, Suresh Kumar^{a,*}, Meena Kumari^b, Manishita Rani^a, Neera Raghav^a, Pranab Kumar Sahoo^c, Sushmita Ghosh^c, Sutapa Mahata^c, Vilas D. Nasare^c

^a Department of Chemistry, Kurukshetra University, Kurukshetra, Haryana, 136119, India

^b Department of Chemistry, Govt. College for Women, Badhra, Charkhi Dadri, Haryana, 127306, India

^c Department of Pathology and Cancer Screening, Chittaranjan National Cancer Institute, 37, S.P. Mukherjee Road, Kolkata, 700026, India

ARTICLE INFO

Keywords:

AGS cell line
Aurone
Cathepsin B
One-pot multicomponent synthesis
1,2,3-Triazole hybrid
Ultrasound

ABSTRACT

Cathepsin B represents a group of lysosomal-encapsulated cellular cysteine proteases, whose upregulation is the critical risk factor for cancer progression and in the regulation of degenerative processes like apoptosis. Therefore, cysteine cathepsins are used as effective biomarkers for cancers and other neurodegenerative disorders and have become specific targets in drug designing. To diversify the pharmacological potential of aurones, a total of eighteen novel 1,2,3-triazole decorated aurone hybrids were synthesized in very good yields, utilizing click chemistry and an ultrasound-assisted one-pot, multicomponent and greener protocol, straight from hydroxy-substituted aurones. These hybrids have been evaluated as useful cathepsin B inhibitors and further explored their *in-vitro* study over AGS cell line using MTT assay. The *in-vitro* cathepsin B inhibitory potential of the synthesized hybrids revealed that the hybrid 3p possesses a very significant inhibition (% inhibition = 85.02) at the concentration of 10^{-6} M compared to the standard aspirin (% inhibition = 48.21 at 10^{-7} M) and curcumin (% inhibition = 57.72 at 10^{-7} M), which further supported by the *in-silico* studies. In addition to this, the cytotoxic assay of the hybrids showed that the hybrids 3d (IC_{50} = 24 μ M), 3i (IC_{50} \approx 24 μ M), and 3m (IC_{50} = 26 μ M) exhibit comparable anti-tumor activity against AGS cell line than the standard, oxaliplatin (IC_{50} = 29 μ M). The results suggested that the reported novel aurone-1,2,3-triazole hybrids are a promising candidate for the development of new cathepsin B inhibitors as well as anti-cancer leads.

1. Introduction

Cancer is a group of appealing diseases which generally arise due to the transformation of healthy cells into tumor cells. This transformation generally occurs due to altered cellular repair mechanism controlled by genetic factors in response to exposure towards certain carcinogens (biological, chemical, or physical). In spite of significant developments in methods of cancer diagnosis and treatment (Radiotherapy/Chemotherapy/Surgery etc.) in last few years, cancer is still a leading cause of death worldwide. The major hurdles in cancer treatment include detection in later stages, resistance towards various drugs used in chemotherapy, and ability to differentiate between normal and cancerous cells. The researchers across the world are continuously working for the design and development of newer cytotoxic agents with magnified efficacy,

minimal toxicity and for effective cancer treatment by following diverse approaches. One of these approaches involve designing of therapeutic molecules which target various enzymes like cysteine proteases and their receptors thereby suppressing their activity [1,2]. These proteases are known to play significant role in the progression of various oncogenic processes [3], and various other tissue degenerative diseases [4,5], like, rheumatoid arthritis, osteoarthritis, inflammation [6], Alzheimer's [7], multiple sclerosis, and pancreatitis [8], etc. in humans. The inhibition of cathepsin proteases is further necessitated for restricting their role in development of both intrinsic and acquired resistance towards chemotherapy and radiation therapy, used in cancer treatment [9]. Cathepsin B is one such ubiquitously expressed cysteine protease whose over-expression is correlated with metastatic and invasive proliferation of cancer [10,11]. Recent data have shown that the progression of some

* Corresponding author. Department of Chemistry, Kurukshetra University, Kurukshetra, Haryana, India.
E-mail address: suresh_dua47@rediffmail.com (S. Kumar).

<https://doi.org/10.1016/j.crgsc.2022.100295>

Received 30 December 2021; Received in revised form 11 February 2022; Accepted 26 February 2022


Available online 2 March 2022

2666-0865/© 2022 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

- medium, *Sustain. Chem. Pharm.* 19 (2021) 100358, <https://doi.org/10.1016/j.scp.2020.100358>.
- [65] M. Boominathan, N. Pugazhenthi, M. Nagaraj, S. Mathanarasimhan, S. Murugesan, N. Bhuvanesh, Nanoporous silica-supported gold nanoparticles-catalyzed green synthesis of 1, 2, 3-triazole in aqueous medium, *ACS Sustainable Chem. Eng.* 1 (2013) 1405–1411, <https://doi.org/10.1021/sc400167a>.
- [66] I. Murthy, R. Sreenivasulu, G. Alluraj, R.R. Raju, Design, synthesis, and anticancer activity of 1, 2, 3-triazole linked 1, 2-lactazole imidazole [4, 5-b] pyridine derivatives, *rucc. J. Gen Chem.* 89 (2019) 1718–1723, <https://doi.org/10.1178/810700019219050270>.
- [67] P. Nouraei, S. Moradi Dehaghi, A. Foroumadi, *Quinazolin-1, 2, 3-triazole hybrid derivatives: green synthesis and DFT calculations*, *Synth Commun* 49 (2019) 386–394, <https://doi.org/10.1080/00907179.2019.1557486>.
- [68] M. Saredi, M. Mohammadi Khanpooshani, F. Pourabbas, N. Rezakhalil, R. Ghadimi, S. Iranparast, M.A. Faramarzi, F. Bandarian, E.N. Fatahali, M. Safari, Design and synthesis of novel quinoxaline-1, 2, 3-triazole hybrid as new anti-diabetic agents: in vitro a glucosidase inhibition, kinetic, and docking study, *Bioorg. Chem.* 83 (2019) 161–169, <https://doi.org/10.1016/j.bioorg.2019.101773>.
- [69] M. Eftekhari, F. Razaqi, Synthesis, characterization and first application of Graphene oxide functionalized Cu (II) complex for the synthesis of 1, 2, 3-triazole derivatives, *Polycycl. Aromat Comp.* (2021) 1–13, <https://doi.org/10.1080/10406638.2021.1913195>.
- [70] J. Anterberry, S. Mokenapelli, G. Thaleri, Facial synthesis of novel 3-(2-methylbenzofuran-3-yl) 5-(4-(phenoxymethyl)-1H-1, 2, 3-triazole-1-yl) methyl)-1, 2, 4-oxadiazole derivatives, *Synth Commun* 51 (2021) 1417–1424, <https://doi.org/10.1080/00397911.2021.1984851>.
- [71] M.A. Alamehadi, A. Aljafari, S.Y. Alragi, I. Ali, N. Rezki, M.R. Aoud, M. Hagar, Design, synthesis, DNA binding, modelling, anti-cancer studies and DFT calculations of Schiff bases tethering benzothiazole-1, 2, 3-triazole conjugates, *J. Mol. Struct.* 1225 (2021) 129148, <https://doi.org/10.1016/j.molstruc.2020.129148>.
- [72] H.K. Siuki, P.G. Karger, G. Bagherzade, Upgraded valorization of copper (II) immobilization onto schiff base-functionalized pectin grafted magnetic nanospheres: a thermally stable, reusable, and efficient nanocatalyst for the green synthesis of 1, 4-disubstituted pyrazole in green water 2 (2021) 3, <https://doi.org/10.1016/j.ces.2020.100358>.
- [73] M. Khatami, M. Shoghi, S. Garg, B. Kave, S. Jangari, I. Shafiei, Ion exchange: a useful tool for separation and simultaneous purification of lysosomal enzyme proteinases, cathepsins B, H and L, *Int. J. Pharm.* 5 (2015) 2548–2558, <https://doi.org/10.1016/j.ijpharm.2015.05.037>.
- [74] S. Khatami, M. Shoghi, Synthesis, purification and docking studies of bisphthalonitrile based quinoxaline-1, 2, 3-triazole and quinoxaline-1, 2, 3-triazole derivatives as novel inhibitors of cathepsins B and cathepsin H, *Int. J. Pharmaceut.* 561 (2014) 38–39, <https://doi.org/10.1016/j.ijpharm.2014.01.018>.
- [75] I. Shafiei, S. Khatami, Comparison of inhibition of mammalian cathepsins B, cathepsin H, and phosphatase and alkaline phosphatase: a correlation with pharmacological activities, *Mol. Chem. Res.* 23 (2014) 2847–2853, <https://doi.org/10.1002/chem.201400072>.
- [76] S. Khatami, S. Garg, S. Kave, S. Shoghi, Synthesis, antimicrobial evaluation, and in silico studies of quinoxaline-1, 2, 3-triazole molecular hybrids, *Mol. Divers.* 25 (2021) 2255–2278, <https://doi.org/10.1007/s11033-020-10131-3>.
- [77] M. Jafari, S. Kave, S. Khatami, S. Garg, S. Kave, S. Shoghi, S. Khatami, M. Akhbari, Synthesis, QSAR and antimicrobial evaluation of 1,2,3-triazole-5-ylidene derived from diamethyl hydrazine derivatives, *Int. J. Med. Chem.* 53 (2015) 2486–2494, <https://doi.org/10.1155/2015/2486>.
- [78] S. Khatami, S. Kave, S. Khatami, S. Garg, S. Kave, S. Shoghi, S. Khatami, Synthesis and antimicrobial activity of 2-substituted 1,2,3-triazole-5-ylidene derivatives, *Molecules* 23 (2018) 1882, <https://doi.org/10.3390/molecules23081882>.
- [79] M.S. Lomonova, D. Serebrennik, R.T. Mikhlin, E.S. Kostin, The total synthesis of an unusual natural from *Ulmus laevis* leaves: structure and discovery of anticancer agents, *Bioorg. Med. Chem. Lett.* 13 (2002) 3759–3763, <https://doi.org/10.1056/b.2002.129148>.

NEW Current Research In Green And Sustainable Chemistry

Overview



I. Basic Journal Info

Country



Netherlands (<https://www.scijournal.org/country/netherlands>)


Journal ISSN: 26660865

Publisher: Elsevier BV (<https://www.scijournal.org/publishers/elsevier-bv>)

History: 2020-2021

Journal Homepage: Link (NA)

How to Get Published:



Novel 1,2,3-triazole-aurone hybrids as cathepsin B inhibitors: One-pot synthesis, anti-proliferative, and drug modeling studies

Bhavna Saroha^a, Gourav Kumar^a, Suresh Kumar^{a,*}, Meena Kumari^b, Manishita Rani^a, Neera Raghav^a, Pranab Kumar Sahoo^c, Sushmita Ghosh^c, Sutapa Mahata^c, Vilas D. Nasare^c

^a Department of Chemistry, Kurukshetra University, Kurukshetra, Haryana, India, 136119

^b Department of Chemistry, Govt. College for Women, Badhra, Charkhi Dadri, Haryana, India, 127306

^c Department of Pathology and Cancer Screening, Chittaranjan National Cancer Institute, 37, S.P. Mukherjee Road, Kolkata, India, 700026

ARTICLE INFO

Keywords:

Aurones
Anti-proliferative activity
Cathepsin B inhibitors
One-pot synthesis
1,2,3-Triazole

ABSTRACT

In the present study, we have designed a series of novel 1,2,3-triazole-aurone hybrids, which were synthesized by a one-pot click reaction between a terminal alkyne and an *in-situ* generated azide, and evaluated as potential cathepsin B inhibitors. The structures of the synthesized compounds were confirmed by various spectroscopic techniques (¹H & ¹³C NMR and HRMS data). The *in-vitro* cathepsin B inhibitory study of the synthesized compounds showed that the hybrids named 3o, 3p, and 3r are the excellent inhibitors with percentage inhibition of 83.05–87.16% at the concentration of 10⁻⁵ M, in comparison to standards, Aspirin (48.21% at 10⁻⁷ M), and Curcumin (52.27% at 10⁻⁷ M). The docking study was also in support of the *in-vitro* assay. Furthermore, the anti-proliferative impact of these synthesized compounds was also evaluated against the gastric adenocarcinoma cell line (AGS cell line). The *in-vitro* MTT assay suggested that out of all the synthesized hybrids, compounds 3a with an IC₅₀ value of 16 μM and 3b with an IC₅₀ value of 11 μM possessed maximum cytotoxic activity against the AGS cell line in comparison to the reference, Oxaliplatin (IC₅₀ = 29 μM).

1. Introduction

Cathepsin B is a ubiquitously expressed mammalian lysosomal peptidase that possesses both carboxypeptidase and endopeptidase activities. The carboxypeptidase activities of the enzyme play diverse roles in various physiological processes like protein processing, wound healing and apoptosis [1], etc. In contrast, the endopeptidase activities like degradation of the extracellular matrix are extremely up-regulated in several pathological conditions like cancer [2,3], inflammation [4], parasitic infection, and neurodegeneration [5,6], etc. The altered trafficking of cathepsin B cause increased proteolytic activity, which promotes the invasion and metastasis of tumor cells, thereby leading to malignant progression and poor patient prognosis.

Thus, to avoid the extreme up-regulation of these cathepsin B enzyme molecules, their activity is regulated at every level, starting from their biosynthesis via compartmentalization into lysosomes, controlled activation of their pro-enzyme forms, and by their endogenous protein inhibitors like cystatins [7]. In addition to these endogenous regulators, various exogenous protein inhibitors of cathepsin B have been designed either by extraction from natural sources or by synthetic preparatory

methods. The analysis and interpretation of x-ray crystal structures of cathepsin B complex with its inhibitors provide further insights into designing various new small molecular weight cathepsin B inhibitors. Our research group is also working actively in the field of medicinal chemistry to design and develop novel heterocyclics as inhibitors of cathepsin B for the treatment of several diseases, like neural disorders, cell proliferation, and various others.

In this process, we have achieved a lead in developing some novel aurone derivatives as potent anti-cancer [8] and cathepsin B inhibitors [9]. Aurones (benzofuranones) are privileged structures in medicinal chemistry depicting a wide range of biological activities [10–13], like anti-cancer [14,15], anti-alzheimer [16,17], anti-inflammatory [18], and various other activities. The pharmacological profile of these aurones can further be tuned by making various other structural modifications [19, 20]. Besides aurones; several other heterocyclic skeletons depicting cathepsin B inhibition and significant anti-cancer activity have been developed, including triazoles [21], chalcones [22], and flavones [23] (Fig. 1) by various research groups. Based on our previous results, we aimed to develop the hybrid compounds of aurones with other biologically active heterocyclic skeletons as novel therapeutic molecules with

* Corresponding author. Department of Chemistry, Kurukshetra University, Kurukshetra, Haryana, India.

E-mail address: suresh_dua47@rediffmail.com (S. Kumar).

<https://doi.org/10.1016/j.ejmc.2022.100056>

Received 29 December 2021; Received in revised form 23 April 2022; Accepted 1 May 2022

Available online 6 May 2022

2772-4174/© 2022 The Authors. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

- [4] A. Lang, D. Horler, A. Balci, The relative importance of cysteine peptidases in osteoarthritis, *J. Rheumatol.* 27 (8) (2000) 1970–1979.
- [5] U. Felton, B. Kessler, W. Mothes, H.H. Goebel, H.L. Ploegh, R.T. Bronson, B.R. Olsen, Neuronal loss and brain atrophy in mice lacking cathepsins B and L, *Proc. Natl. Acad. Sci. Unit. States Am.* 99 (12) (2002) 7883–7888.
- [6] V. Hook, M. Yoon, C. Mosler, G. Ito, S. Podvin, B.P. Head, R. Rissman, A.J. O'Donoghue, G. Hook, Cathepsin B in neurodegeneration of Alzheimer's disease, traumatic brain injury, and related brain disorders, *Biochim. Biophys. Acta Protein Proteomics* (2020), 140428.
- [7] R. Frilan, S. Gobec, Inhibitors of cathepsin B, *Curr. Med. Chem.* 13 (19) (2006) 2309–2327.
- [8] S. Kumar, E. Lathwal, G. Kumar, B. Saroha, S. Kumar, S. Mahata, P.K. Sahoo, V.D. Nasare, Synthesis of pyrazole based novel aurone analogs and their cytotoxic activity against MCF-7 cell line, *Chem. Data Collect.* 30 (2020), 100559.
- [9] B. Saroha, G. Kumar, E. Lathwal, S. Kumar, M. Kumari, N. Mor, N. Raghav, S. Kumar, Synthesis of propynyloxy substituted some novel aurones as potent cathepsin B inhibitors, *Chem. Data Collect.* 31 (2021), 100630.
- [10] G. Kumar, E. Lathwal, B. Saroha, S. Kumar, S. Kumar, N.S. Chauhan, T. Kumar, Synthesis and biological evaluation of quinoline-based novel aurones, *ChemistrySelect* 5 (12) (2020) 3539–3543.
- [11] V.R. Pereira, L.S. da Silveira, A.C. Meigarda, J.J.A. Júnior, O.O.Z. da Silva, F.B. Miguel, M.P. Silva, A.d.C. Almeida, D. da Silva Torres, P.d.F. Pinto, Antischistosomal properties of aurone derivatives against juvenile and adult worms of *Schistosoma mansoni*, *Acta Trop.* 213 (2021), 105741.
- [12] S. Paidakula, S. Nerella, R. Vadde, A. Kamal, S. Kankala, Design and synthesis of 4p-Acetamidobenzofuranone-podophyllotoxin hybrids and their anti-cancer evaluation, *Bioorg. Med. Chem.* 29 (16) (2019) 2153–2156.
- [13] A.S. Chintakindi, D.J. Gohil, A.S. Chowdhary, M.A. Kanyalkar, Design, synthesis and biological evaluation of substituted flavones and aurones as potential anti-influenza agents, *Bioorg. Med. Chem.* 28 (1) (2020), 115191.
- [14] A.-H. Liu, L.-H. Liu, C.-Y. Liu, Anticancer potential of (2 Z)-2-[(4'-Hydroxyphenyl) methylene]-6-methoxy-7-prenyl-3 (2 H)-Benzofuran, a new aurone from *Psoralea corylifolia*, *Nat. Prod. Commun.* 15 (1) (2020), 1934578X19899505.
- [15] N. Verma, K. Chauhan, S. Kumar, In benzofuran-3 (2H)-ones derivatives: synthesis, docking and evaluation of their in vitro anticancer studies, in: *Proceedings of International Conference on Trends in Computational and Cognitive Engineering*, Springer, 2021, pp. 317–325.
- [16] K.-F. Liew, K.-I. Chan, C.-Y. Lee, Blood-brain barrier permeable anticholinesterase aurones: synthesis, structure-activity relationship, and drug-like properties, *Eur. J. Med. Chem.* 94 (2015) 195–210.
- [17] Y. Li, X. Qiang, L. Luo, Y. Li, G. Xiao, Z. Tan, Y. Deng, Synthesis and evaluation of 4-hydroxyl aurone derivatives as multifunctional agents for the treatment of Alzheimer's disease, *Bioorg. Med. Chem.* 24 (10) (2016) 2342–2351.
- [18] B.P. Bandgar, S.A. Patil, B.L. Korbadi, S.C. Biradar, S.N. Nile, C.N. Khobragade, Synthesis and biological evaluation of a novel series of 2, 2-bisaminomethylated aurone analogues as anti-inflammatory and antimicrobial agents, *Eur. J. Med. Chem.* 45 (7) (2010) 3223–3227.
- [19] A. Meguellati, A. Ahmed-Belkacem, W. Yi, R. Haudecoeur, M. Crouillere, R. Brillet, J.-M. Pawlowsky, A. Boumendjel, M. Peuchmaur, B-ring modified aurones as promising allosteric inhibitors of hepatitis C virus RNA dependent RNA polymerase, *Eur. J. Med. Chem.* 80 (2014) 579–592.
- [20] H. Olleik, E. Courvoisier-Dezord, J. Perrier, S. Yahiaoui, B. Roulier, B. Peres, A. Boumendjel, A. Hijazi, E. Baydoun, J. Raymond, M. Maresca, R. Haudecoeur, Aurone derivatives as promising antibacterial agents against resistant Gram-positive pathogens, *Eur. J. Med. Chem.* 165 (2019) 133–141.
- [21] N. Raghav, M. Singh, Acyl hydrazides and triazoles as novel inhibitors of mammalian cathepsin B and cathepsin H, *Eur. J. Med. Chem.* 77 (2014) 231–242.
- [22] D.H. Soliman, A.M. Farrag, O. Omsan, Design, synthesis and in-silico studies of novel chalcones as anti-prostate cancer and cathepsin B inhibitors, *J. Appl. Pharmaceut. Sci.* 7 (2017) (07), 010–020.
- [23] L.M. de Nuyvis, C.C. de Azeiteiro, I.F. Ferreira, T.A. Ribeiro, P.T. Sousa Jr., M.J. Jacinto, M.G. de Carvalho, W.A. Judice, L.O. Jesus, A.A. de Sousa, 4'-Hydroxy-6, 7-methylenedioxy-3-methoxyflavone: a novel flavonoid from *Dulacina egleri* with potential inhibitory activity against cathepsins B and L, *Phytorespon.* 132 (2019) 26–29.
- [24] B. Yu, P.-P. Qi, X.-J. Shi, R. Huang, H. Guo, Y.-C. Zheng, D.-Q. Yu, H.-M. Lin, Efficient synthesis of new antiproliferative steroid hybrids using the molecular hybridization approach, *Eur. J. Med. Chem.* 117 (2016) 241–255.
- [25] N. Nehra, R.K. Tirol, D.G. Vikas, K. Lal, Synthesis, antifungal studies, molecular docking, ADME and DNA interaction studies of 4-hydroxyphenyl benzothiazole linked 1, 2, 3-triazoles, *J. Mol. Struct.* 1245 (2021), 131012.
- [26] B. Saroha, G. Kumar, R. Kumar, M. Kumari, S. Kumar, A mini-review of 1, 2, 3-Triazole hybrids with O-Heterocycles as leads in medicinal chemistry, *Chem. Biol. Drug Des.* (2021), <https://doi.org/10.1111/cbdd.13966>.
- [27] D. Dheer, V. Singh, R. Shankar, Medicinal attributes of 1, 2, 3-triazoles: current developments, *Bioorg. Chem.* 71 (2017) 30–54.
- [28] O. Ivashkevich, V.E. Marulis, P. Gaponik, G. Sukhanov, J. Filipponi, A. Sukhanova, Quantum-chemical investigation of certain physicochemical properties of 6-nitro-1, 2, 3-triazole and N-alkyl-4 (5)-nitro-1, 2, 3-triazoles, *Chem. Heterocycl. Compd.* 44 (12) (2008) 1472–1482.
- [29] H. Li, R. Aneja, I. Chaiken, Click chemistry in peptide-based drug design, *Molecules* 18 (8) (2013) 9797–9817.
- [30] C. Gao, L. Chang, Z. Xu, X.-F. Yan, C. Ding, F. Zhao, X. Wu, L.-S. Feng, Recent advances of tetrazole derivatives as potential anti-tubercular and anti-malarial agents, *Eur. J. Med. Chem.* 163 (2019) 404–412.
- [31] A. Djemoui, A. Naoun, M.R. Ouahani, D. Djemoui, S. Labenne, M.B. Labrech, I. Boukenna, H.M. Albuquerque, L. Soher, D.H. Rocha, A step-by-step synthesis of triazole-benzimidazole-chalcone hybrids: anticancer activity in human cells, *J. Mol. Struct.* 1204 (2020), 127487.
- [32] A.B. Kasetti, I. Singhvi, R. Nagasuri, R.R. Bhandare, A.B. Shaik, Thiazole-Chalcone hybrids as prospective antitubercular and antiproliferative agents: design, synthesis, biological, molecular docking studies and in silico ADME evaluation, *Molecules* 26 (10) (2021) 2847.
- [33] T. Sowjanya, Y.J. Rao, N. Murthy, Synthesis and antiproliferative activity of new 1, 2, 3-triazole/flavone hybrid heterocycles against human cancer cell lines, *Russ. J. Gen. Chem.* 87 (8) (2017) 1864–1871.
- [34] A. Rastegari, H. Nadri, M. Mahdavi, A. Moradi, S.S. Mirfathi, N. Ebrahimi, F.H. Moghadam, R. Larijani, T. Akbarzadeh, M. Soerch, Design, synthesis and anti-Alzheimer's activity of novel 1, 2, 3-triazole-chromone carboxamide derivatives, *Bioorg. Chem.* 83 (2019) 391–401.
- [35] M. Yazdani, N. Ebrahimi, R. Radhi, M. Khoshneviszadeh, A. Irani, O. Firuzi, S. 6-Diphenyl triazine-thio methyl triazole hybrid as a new Alzheimer's disease modulating agents, *Mol. Divers.* 24 (3) (2020) 641–654.
- [36] N. Raghav, R. Kam, Chalcones, semicarbazones and pyrazolines as inhibitors of cathepsins B, H and L, *Int. J. Biol. Macromol.* 80 (2015) 710–724.



Sources

ISSN

ISSN: 27724174 x

i Improved CiteScore

We have updated the CiteScore methodology to ensure a more robust, stable and comprehensive metric which provides an indication of research impact, earlier. The updated methodology will be applied to the calculation of CiteScore, as well as retroactively for all previous CiteScore years (ie. 2018, 2017, 2016...). The previous CiteScore values have been removed and are no longer available.

[View CiteScore methodology.](#)

Filter refine list

1 result

☐ All [Download Scopus Source List](#) [Learn more about Scopus Source List](#)

2022

View metrics for year:

Display options

☐ Display only Open Access journals

Counts for 4-year timeframe

☐ No minimum selected☐ Minimum citations

Source title ↓

CiteScore ↓ Highest percentile ↓

Citations 2019-22 ↓ Documents 2019-22 ↓ % Cited ↓

☐ 1 European Journal of Medicinal Chemistry Reports *Open Access*

24% 49/64

47

86

114

Chemistry (miscellaneous)

Polyglycerol-based hydrogels and nanogels: from synthesis to applications

Meena Kumari^{1,2}, Suchita Prasad^{1,3}, Ljiljana Fruk^{*1} & Badri Parshad^{**1}

¹Department of Chemical Engineering & Biotechnology, University of Cambridge, Cambridge, CB3 0AS, UK

²Department of Chemistry, Government College for Women, Badhra, Ch. Dadri, Haryana 127308, India

³Department of Chemistry, University of Delhi, Delhi 110007, India

*Author for correspondence: lf389@cam.ac.uk

**Author for correspondence: bp448@cam.ac.uk

†Authors contributed equally

Hydrogels and nanogels have emerged as promising materials for biomedical applications owing to their large surface area and tunable mechanical and chemical properties. Their large surface area is well suited for bioconjugation, whilst the interior porous network can be utilized for the transport of valuable biomolecules. The use of biocompatible hydrophilic building blocks/linkers for the preparation of hydrogels and nanogels not only avoids undesired side effects within the biological system, but also retains high water content, thereby creating an environment which is very similar to extracellular matrix. Their tunable multivalency and hydrophilicity and excellent biocompatibility, together with ease of functionalization, makes polyglycerol macromonomers well suited for synthesizing cross-linked networks that can be used as extracellular matrix mimics. Here we provide an overview of the synthesis of polyglycerol-based hydrogels and nanogels for various biomedical applications.

First draft submitted: 25 June 2020; Accepted for publication: 1 December 2020; Published online: 6 January 2021

Keywords: chemical cross-linking • hydrogel • nanogel • polyglycerol

Hydrogels are 3D cross-linked networks capable of retaining a large amount of water. Due to the possibility to tune their biocompatibility, molecular flexibility, stiffness and binding affinity for biomolecules by various strategies, they have been widely explored as mimics of extracellular matrix. Such matrix is a heterogeneous structure unique to each tissue that contains the physical, biochemical and biomechanical components needed for cell proliferation and communication. The development of extracellular matrices with the desired properties is one of the most important steps in tissue engineering and has been benefited largely from advances in the design of different classes of cross-linked hydrogels [1]. Besides tissue engineering, hydrogels have made a significant contribution in the fields of drug delivery, pathogen inhibition, biomolecule immobilization and the production of hygiene products, wound dressings and contact lenses [2]. The conjugation of suitable ligands on the surface of cross-linked networks further facilitates the delivery of encapsulated guests to the target site as well as improves their affinity to bind with specific pathogens.

The current field of hydrogel research commenced with the preparation of a polyhydroxyethylmethacrylate (pHEMA) hydrogel by Wichterle and Lim in 1960, when these materials were utilized for contact lens applications [3]. The first nanogel was prepared three decades later by self-aggregation of cholesterol-bearing polysaccharides in water that resulted in physically cross-linked nanosized structures [4]. Nanogels are fabricated using various methodologies, such as inverse miniemulsion, inverse nanoprecipitation and microfluidics, and with various cross-linking species. They have received significant attention in recent years owing to their nanosized features and easily tailored chemical and physical properties and have been used as space-filling agents, delivery vehicles for bioactive molecules [5–8], biosensors [9–12], bioimaging [13–16] and antifouling agents [17,18], and also as tissue engineering elements [19–22].

Several excellent review articles have been published concerning hydrogel and nanogel design [23–26], but here we will focus on polyglycerol (PG)-based systems, which have been extensively explored in the past decade.

newlands
press

- 60 Zabihi F, Wiczczyk S, Dimde M, Hedtrich S, Börner HG, Haag R. Intradermal drug delivery by nanogel-peptide conjugates: specific and efficient transport of temoporfin. *J. Control. Release* 242, 35–41 (2016).
- 61 Dimde M, Neumann F, Reisbeck P *et al.* Defined pH-sensitive nanogels as gene delivery platform for siRNA mediated *in vitro* gene silencing. *Biomater. Sci.* 5(11), 2328–2336 (2017).
- 62 Dimde M, Söhle FF, Wycisk V *et al.* Synthesis and validation of functional nanogels as pH-sensors in the hair follicle. *Macromol. Biosci.* 17(10), 1600505 (2017).
- 63 Obst K, Yealland G, Balzus B *et al.* Protein corona formation on colloidal polymeric nanoparticles and polymeric nanogels: impact on cellular uptake, toxicity, immunogenicity, and drug release properties. *Biomacromolecules* 18(6), 1762–1771 (2017).
- 64 Zhang X, Achazi K, Steinhilber D, Kratz F, Dornedde J, Haag R. A facile approach for dual-responsive prodrug nanogels based on dendritic polyglycerols with minimal leaching. *J. Control. Release* 174, 209–216 (2014).
- 65 Zhang X, Achazi K, Haag R. Boronate cross-linked ATP- and pH-responsive nanogels for intracellular delivery of anticancer drugs. *Adv. Healthcare Mater.* 4(4), 585–592 (2015).
- 66 Zhang X, Zhang K, Haag R. Multi-stage, charge convertible, stimuli-responsive nanogels for therapeutic protein delivery. *Biomater. Sci.* 3(11), 1487–1496 (2015).
- 67 Steinhilber D, Sisson AL, Mangoldt D, Welker P, Licha K, Haag R. Synthesis, reductive cleavage, and cellular interaction studies of biodegradable, polyglycerol nanogels. *Adv. Funct. Mater.* 20(23), 4133–4138 (2010).
- 68 Zhou H, Steinhilber D, Schlaad H, Sisson AL, Haag R. Glycerol based polyether-nanogels with tunable properties via acid-catalyzed epoxide-opening in miniemulsion. *React. Funct. Polym.* 71(3), 356–361 (2011).
- 69 Hellmund M, Zhou H, Samsonova O, Welker P, Kissel T, Haag R. Functionalized polyglycerol amine nanogels as nanocarriers for DNA. *Macromol. Biosci.* 14(9), 1215–1221 (2014).
- 70 Gao L, Zabihi F, Ehrmann S, Hedtrich S, Haag R. Supramolecular nanogels fabricated via host-guest molecular recognition as penetration enhancer for dermal drug delivery. *J. Control. Release* 300, 64–72 (2019).
- **Produces nanogel via host-guest interactions as cross-linking strategy.**
- 71 Oehrl A, Schütz S, Haag R. Systematic screening of different polyglycerol-based dienophile macromonomers for efficient nanogel formation through IEDDA inverse nanoprecipitation. *Macromol. Rapid Commun.* 41(1), 1900510 (2020).
- 72 Stolz L, Vigovskaya A, Barner-Kowollik C, Fruk L. A self-reporting tetrazole-based linker for the biofunctionalization of gold nanorods. *Chem. Eur. J.* 21(41), 14309–14313 (2015).
- 73 Stolz L, Quick AS, Abt D *et al.* Photo-induced surface encoding of gold nanoparticles. *Chem. Commun.* 51(16), 3363–3366 (2015).

-NEW Future Medicinal Chemistry

Overview

Impact Factor

3.808




H Index

76



Impact Factor

3.643



I. Basic Journal Info

Country



United Kingdom (<https://www.scijournal.org/country/united-kingdom>)

Journal ISSN: 17568919, 17568927

Publisher: Future Medicine Ltd. (<https://www.scijournal.org/publishers/future-medicine-ltd>)

History: 2009-ongoing

Journal Homepage: [Link \(NA\)](#)

How to Get Published:



Enzymatic synthesis of glycerol, azido-glycerol and azido-triglycerol based amphiphilic copolymers and their relevance as nanocarriers: A review

Badri Parshad^{a,*}, Meena Kumari^b, Vinod Khatri^c, Rajeshwari Rajeshwari^d, Yuanwei Pan^e,
Atul K. Sharma^f, Ishtiaq Ahmed^a

^a Department of Chemical Engineering and Biotechnology, University of Cambridge, Cambridge CB3 0AS, UK

^b Department of Chemistry, Government College for Women Badhra, Ch. Dadri, Haryana, India

^c Department of Chemistry, T.D.L. Govt. College for Women, Murthal, Sonapat, Haryana, India

^d Department of Chemistry, Umeå University, 90187 Umeå, Sweden

^e Institut für Chemie und Biochemie, Freie Universität Berlin, Takustraße 3, 14195 Berlin, Germany

^f Department of Chemistry, University of Delhi, Delhi 110007, India

ARTICLE INFO

Keywords:

Glycerol
Azido-glycerol
Copolymer
Amphiphilic
Self-assembly
Nanocarrier

ABSTRACT

Amphiphilic polymeric nanocarriers have attained immense attention for transporting drugs and other bioactive species to the living systems owing to their high loading capacity and efficient internalization. To avoid the side-effects of undesired interactions within the biological system, the use of biocompatible building blocks is most crucial in preparing polymeric amphiphiles. The excellent biocompatibility and multivalency of both glycerol and triglycerol make them suitable monomers to construct macromolecular skeletons. Besides these, easy availability and remarkable aqueous solubility further enlarge their use in synthetic chemistry and give ample possibilities for creating fascinating entities for practical applications. The conversion of glycerol into azido-glycerol and azido-triglycerol further helped in differential functionalization of their polymers with various hydrophobic and hydrophilic groups in different ratios thereby assisting in tuning the amphiphilicity of resulting functionalized polymers. Herein, we review the enzymatic synthesis of glycerol, azido-glycerol and azido-triglycerol based amphiphilic polymeric architectures as nanocarriers for various bio-active species. Enzymatically synthesized linear copolymers synthesized by selective esterification of primary hydroxyl groups of glycerol and its derivatives with suitable diacids/diesters are explored in this review.

1. Introduction

Current advances in the field of drug delivery have revealed the substantial importance of amphiphilic polymeric architectures derived from bio-based monomers as proficient nanocarriers for the transport of bioactive molecules to the target site within the biological system [1–6]. To confront the challenges faced by most of the drugs which include low water solubility, high drug doses and a short lifetime in the bloodstream

[7–11], researchers across the world have focused to improve the pharmacokinetics of the drugs by developing amphiphilic polymeric nanocarriers produced from biocompatible starting materials. Among the huge list of various starting monomeric building blocks, glycerol, possessing excellent biocompatibility, is highly successful in catching the eye of researchers [12–15]. Since glycerol is obtained as a by-product in huge amounts (~10% w/w) from bio-diesel productions, it is easily available at a very low cost across the world [16]. It is also

Abbreviations: ATRP, Atom transfer radical polymerization; CAC, Critical aggregation concentration; CAL B, *Candida antarctica* lipase B; CLSM, Confocal laser scanning microscopy; CMC, Critical micelle concentration; CNT, Carbon nanotube; CuAAC, Copper-catalyzed azide-alkyne cycloaddition; DAPI, 4',6-Diamidino-2-Phenylindole; DBTO, Dibutyltin oxide; DCC, N,N'-Dicyclohexylcarbodiimide; DIPEA, N,N-Diisopropylethylamine; DLS, Dynamic light scattering; DMF, Dimethylformamide; FACS, Fluorescence-assisted cell sorting; FDA, Food and drug administration; FTIR, Fourier transform infrared; G1, First generation; GPC, Gel permeable chromatography; GRAS, Generally recognized as safe; hPG, hyperbranched polyglycerol; MALDI-TOF, Matrix-assisted laser desorption/ionization-Time of flight; MTX, Methotrexate; N435, Novozym 435; NMR, Nuclear magnetic resonance; PBI, Perylene bisimide; PBS, Phosphate buffer saline; PEG, Polyethylene glycol; PDI, polydispersity index; PG, Polyglycerol; SWCNT, Single-walled carbon nanotube; TEM, Transmission electron microscopy; THF, Tetrahydrofuran; 2-MeTHF, 2-Methyltetrahydrofuran.

* Corresponding author.

E-mail address: bp448@cam.ac.uk (B. Parshad).

<https://doi.org/10.1016/j.eurpolymj.2021.110690>

Received 24 May 2021; Received in revised form 30 July 2021; Accepted 3 August 2021

Available online 5 August 2021

0014-3057/© 2021 Elsevier Ltd. All rights reserved.

- [108] W.-X. Wu, N. Wang, B.-Y. Liu, Q.-F. Deng, X.-Q. Yu, Lipase-catalyzed syntheses of azido-functionalized aliphatic polyesters towards acid-degradable amphiphilic graft copolymers, *Soft Matter* 10 (2014) 1199–1213.
- [109] M. Wyszogrodzka, R. Haag, A convergent approach to biocompatible polyglycerol “Click” dendrons for the synthesis of modular core-shell architectures and their transport behavior, *Chem. – A Eur. J.* 14 (2008) 9202–9214.
- [110] M. Wyszogrodzka, K. Mows, S. Kamlage, J. Wodzińska, R. Plietker, R. Haag, New approaches towards monoamino polyglycerol dendrons and dendritic triblock amphiphiles, *Eur. J. Org. Chem.* 2008 (2008) 53–63.
- [111] A. Mittal, A.K. Singh, A. Kumar, K. Parmanand, P. Achazi, S.K. Sharma, R. Haag, Fabrication of oligo-glycerol based hydrolase responsive amphiphilic nanocarriers, *Polym. Adv. Technol.* 31 (2020) 1208–1217.
- [112] S. Kumar, K. Ludwig, B. Schade, H. von Berlepsch, I. Fapp, R. Tyagi, M. Gullia, R. Haag, C. Bortcher, Introducing chirality into nonionic dendritic amphiphiles and studying their supramolecular assembly, *Chem. – Eur. J.* 22 (2016) 5629–5636.
- [113] S. Kumar, K. Achazi, C. Bortcher, K. Licha, R. Haag, S.K. Sharma, Encapsulation and cellular internalization of cyanine dye using amphiphilic dendronized polymers, *Eur. Polym. J.* 69 (2015) 416–428.
- [114] S. Kumar, K. Achazi, K. Licha, P. Manchanda, R. Haag, S.K. Sharma, Chemo-enzymatic synthesis of dendronized polymers for cyanine dye encapsulation, *Adv. Polym. Tech.* 37 (2018) 1797–1805.
- [115] K. Huth, M. Glaeske, K. Achazi, G. Gordeev, S. Kumar, R. Arenal, S.K. Sharma, M. Adeli, A. Setaro, S. Reich, R. Haag, Fluorescent polymer–single-walled carbon nanotube complexes with charged and noncharged dendronized perylene bisimides for biomaging studies, *Small* 14 (2018) 1806796.

NEW European Polymer Journal

Overview



I. Basic Journal Info

Country



United Kingdom (<https://www.scijournal.org/country/united-kingdom>)

Journal ISSN: 143057

Publisher: Elsevier Ltd. (<https://www.scijournal.org/publishers/elsevier-ltd>)

History: 1965-ongoing

Journal Homepage: Link
(http://www.elsevier.com/wps/find/journaldescription.cws_home/294/description#description)

How to Get Published:



Review

A decennary update on diverse heterocycles and their intermediates as privileged scaffolds for cathepsin B inhibition

Bhavna Saroha^a, Gourav Kumar^a, Meena Kumari^b, Ravinder Kaur^a, Neera Raghav^a, Pawan K. Sharma^a, Naveen Kumar^c, Suresh Kumar^{a,*}

^a Department of Chemistry, Kurukshetra University, Kurukshetra, Haryana 136119, India

^b Department of Chemistry, Govt. College for Women, Badli, Charkhi Dadri, Haryana 127308, India

^c Department of Chemistry, Maharshi Dayanand University, Rohtak, Haryana 124001, India

ARTICLE INFO

Keywords:

Cathepsin B (CTSB)

Inhibitors

N-heterocycles

O-heterocycles

SAR

Synthesis

ABSTRACT

The identification of x-ray crystal structure of cathepsin B (CTSB) in the early 90's enabled researchers to embark on a journey to understand and demystify its multiple catalytic mechanisms (endopeptidase/carboxypeptidase/peptidyl-di-peptidase) in diverse physiological processes and their switching into one another under different conditions. The engagement of CTSB in different pathological conditions due to its over-expression further highlighted the enhanced research interest around the domain. The occurrence of over-expressed CTSB in various diseases like Alzheimer's, cancer, arthritis, cardiovascular, etc., and the use of CTSB inhibitors for the treatment of these diseases have established its involvement in different pathological conditions. Such an understanding tempted researchers to design, synthesize, and screen diverse classes of compounds against CTSB. This in turn, helped in understanding their interactions with the active sites of the enzyme. Heterocyclic compounds comprise a very rich and broad class of medicinally important compounds that also hold great potential for CTSB inhibition. This review covers the CTSB inhibition potential of various natural and synthetic heterocyclic scaffolds. Researchers working in the fields of molecular modeling, drug design and development, and enzyme inhibitors can benefit significantly from this review.

1. Introduction

The continuous emergence of newer insights with the ongoing research in the realm of medicinal chemistry facilitates a better understanding of different physiological processes in biological systems, their regulatory pathways, and the association of their malfunctioning with various diseases. The regulation of these physiological processes is done either through enzymes/hormones/receptors, or some other biological molecules. These molecules play a vital role in the proper functioning of the biological system; however, malfunctioning of some of these

regulatory biological molecules is also linked with various diseases. Cathepsin B is one such lysosomal encapsulated cysteine protease enzyme [1–3], which is ubiquitously expressed and involved in several physiological processes [4–7], such as protein degradation and processing, maintaining intracellular homeostatic metabolic activity, tissue remodeling, inflammatory responses against antigens, apoptosis etc. The activity of cathepsin B is controlled by various endogenous inhibitors (Stefins and Cystatins) under normal physiological conditions. The loss of balance between cathepsin B and its inhibitors is marked by the over-expression and unusual localization (cytoplasm/ membrane/

Abbreviations: 1L-1 β , cytokine interleukin-1 β ; A β , β -amyloid; A549, lung cancer cell line; ADME/Tox, Absorption, Distribution, Metabolism, Excretion and Toxicity; APP, Alzheimer's amyloid precursor protein; BANA, N- α -Benzoyl-D, L-arginine- β -naphthylamide hydrochloride; E64d, Aloisistatin (cysteine protease inhibitor); ERG/EWG, Electron releasing groups/Electron withdrawing groups; GI₅₀, concentration causing 50 % cell growth inhibition; HEPG2, liver cancer cell line; HSC-4, human tongue squamous carcinoma cell line; IC₅₀, Half-maximal inhibitory concentration; k₇₇₇, N-[(2S)-1-[[[(E,3S)-1-(benzenesulfonyl)-5-phenylpent-1-en-3-yl]amino]-1-oxo-3-phenylpropan-2-yl]-4-methylpiperazine-1-carboxamide; K_i, inhibition constant; MCF7, breast cancer cell line; MDA-MB-435, metastatic human breast cancer; MTT, (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) tetrazolium; NLRP3, NLR Family Pyrin Domain Containing 3; p62, ubiquitin-binding protein (also known as SQSTM1) involved in autophagy; RI, Relative inhibition; RA, Residual activity; SRB, Sulforhodamine B; SK-LU-1, lung adenocarcinoma cell line; CaSki, HeLa, cervical cancer cell line; DU-145, PC3, prostate cancer cell line.

* Corresponding author.

E-mail address: suresh_dua47@rediffmail.com (S. Kumar).

<https://doi.org/10.1016/j.ijbiomac.2022.10.017>

Received 27 July 2022; Received in revised form 17 September 2022; Accepted 3 October 2022

Available online 8 October 2022

0141-8130/© 2022 Elsevier B.V. All rights reserved.

- [55] S.D. Ramalho, L.R.F. de Sousa, M.C.M. Burger, M.I.S. Lima, M.F.D.G.F. da Silva, J. B. Fernandes, P.C. Vieira, Evaluation of flavonols and derivatives as human cathepsin B inhibitor, *Nat. Prod. Res.* 29 (2015) 2212–2214, <https://doi.org/10.1080/14788419.2014.1092404>.
- [56] N. Suparji, G. Chan, H. Sapili, N. Arshad, L.L. In, K. Awang, N. Hasima Nagoor, Geranylated 4-phenylcoumarins exhibit anticancer effects against human prostate cancer cells through caspase-independent mechanism, *PLoS One* 11 (2016), e0151472, <https://doi.org/10.1371/journal.pone.0151472>.
- [57] L.M.R. de Novais, C.C.O. de Azeiteira, L.F. Ferreira, T.A.N. Ribeiro, P.T. Sousa, M. J. Jacinto, M.G. de Carvalho, W.A.S. Judice, L.O.P. Jesus, A.A. de Souza, H.F. V. Torquato, E.J. Paredes-Gamero, V.C. Silva, 4'-Hydroxy-6,7-methylenedioxy-3-methoxyflavone: a novel flavonoid from *dulacia egleri* with potential inhibitory activity against cathepsins B and L, *Fitoterapia* 132 (2019) 26–29, <https://doi.org/10.1016/j.fitote.2018.08.005>.
- [58] G.O.S. Cunha, A.C.S. Menezes, J.A.D. Silva, A.P. Matos, P.C. Vieira, J.B. Fernandes, M.O.D. Moraes Filho, C.D.O. Pessoa, M.C.D.M. Burgêr, Isolated compounds and semi-synthetic derivatives from *Miconia ferruginata* as inhibitors of cathepsins K and B and their cytotoxic evaluation, *Rev. Virtual Quim.* 12 (2020) 1244–1252, <https://doi.org/10.21577/1984-6835.20200100>.
- [59] V. Caccarini, M. Cuccioloni, Y. Zheng, L. Bonfili, C. Gong, M. Angeletti, P. Mena, D. Del Rio, A.M. Eleuteri, Flavan-3-ol microbial metabolites modulate proteolysis in neuronal cells reducing amyloid-beta (1–42) levels, *Mol. Nutr. Food Res.* 65 (2021) 2100380, <https://doi.org/10.1002/mnfr.202100380>.
- [60] L. Ulčakar, M. Novinec, Inhibition of human cathepsins B and L by caffeic acid and its derivatives, *Biomolecules* 11 (2021) 31, <https://doi.org/10.3390/biom11010031>.
- [61] V. Martichonok, C. Plouffe, A.C. Storer, R. Menard, J.B. Jones, Aziridine analogs of [(trans-Epoxy)succinyl]-L-leucyl]amino]-4-guanidinobutane (E-64) as inhibitors of cysteine proteases, *J. Med. Chem.* 38 (1995) 3078–3085, <https://doi.org/10.1021/jm00016a011>.
- [62] H. Konno, K. Kubo, H. Makabe, E. Toshiro, N. Hinoda, K. Nosaka, K. Akaji, Total synthesis of miraziridine A and identification of its major reaction site for cathepsin B, *Tetrahedron* 63 (2007) 9502–9513, <https://doi.org/10.1016/j.tet.2007.06.082>.
- [63] N.A. Meanwell, Improving drug candidates by design: a focus on physicochemical properties as a means of improving compound disposition and safety, *Chem. Res. Toxicol.* 24 (2011) 1420–1456, <https://doi.org/10.1021/cr200211v>.
- [64] X. Zhang, X. Yang, H. Wang, S. Li, K. Guo, D. Jiang, J. Xiao, D. Liang, Design, synthesis, and structure-activity relationship study of epoxysuccinyl-peptide derivatives as cathepsin B inhibitors, *Biol. Pharm. Bull.* (2017), b17-00075, <https://doi.org/10.1248/bpb.17-00075>.
- [65] T. Van de Walle, M. Briand, A. Mitrović, I. Sostić, S. Gobec, J. Kos, L. Persoons, D. Daelemans, S. De Jonghe, Z. Ubiparip, T. Desmet, K. Van Hecke, S. Mangelinckx, M. D'hooghe, Synthesis of novel nitroxoline analogs with potent cathepsin B exopeptidase inhibitory activity, *ChemMedChem* 15 (2020) 2477–2490, <https://doi.org/10.1002/cmdc.202000402>.
- [66] S. Hartmann, R. Lopez-Cruz, S. Alameh, C.-L.C. Ho, A. Rabideau, B.L. Pentelute, K. A. Bradley, M. Martchenko, Characterization of novel piperidine-based inhibitor of cathepsin B-dependent bacterial toxins and viruses, *ACS Infect. Dis.* 4 (2018) 1235–1245, <https://doi.org/10.1021/acsinfdis.5b00053>.
- [67] Y. Gu, Y. Zhu, G. Deng, S. Liu, Y. Sun, W. Lv, Curcumin analogue AI-44 alleviates MSU-induced gouty arthritis in mice via inhibiting cathepsin B-mediated NLRP3 inflammasome activation, *Int. Immunopharmacol.* 93 (2021), 107375, <https://doi.org/10.1016/j.intimp.2021.107375>.
- [68] L. Cianni, F.D.R. Rocho, V. Bonatto, F.C.P. Martins, J. Lameira, A. Leitão, C. A. Montanari, A. Shamim, Design, synthesis and stepwise optimization of nitrile-based inhibitors of cathepsins B and L, *Bioorg. Med. Chem.* 29 (2021), 115827, <https://doi.org/10.1016/j.bmc.2020.115827>.
- [69] B. Salehi, C. Quispe, I. Chamkhi, N. El Omari, A. Balahbib, J. Sharifi-Rad, A. Bouyahya, M. Akram, M. Iqbal, A.O. Docea, C. Caruntu, G. Leyva-Gómez, A. Dey, M. Martorell, D. Calina, V. López, F. Les, Pharmacological properties of chalcones: a review of preclinical including molecular mechanisms and clinical evidence, *Front. Pharmacol.* 11 (2021), <https://doi.org/10.3389/fphar.2020.592654>.
- [70] A. Rani, A. Anand, K. Kumar, V. Kumar, Recent developments in biological aspects of chalcones: the odyssey continues, *Expert Opin. Drug Dis.* 14 (2019) 249–288, <https://doi.org/10.1080/14746044.2019.1573812>.
- [71] V. Kumar, S. Dhawan, P.S. Girase, P. Awolade, S.R. Shinde, R. Karpoornath, P. Singh, Recent advances in chalcone-based anti-cancer heterocycles: a structural and molecular target perspective, *Curr. Med. Chem.* 28 (33) (2021) 6805–6845, <https://doi.org/10.2174/0929867328666210322102836>.
- [72] A. Owaba, O. Miedegha, R. Oladiran, Chalcones as sythons for heterocyclic compounds: a review, *Int. J. Curr. Res.* 12 (2020) 13672–13681.
- [73] B.H. Tareh, Synthesis, pharmacological activity and uses of chalcone compounds: a review, *Sci. J. Med. Res.* 6 (2022), <https://doi.org/10.37623/sjomr.v06i21.9>.
- [74] M. Rudrapal, J. Khan, A.A.B. Dukhyil, R.M.I.I. Alarousy, E.I. Attah, T. Sharma, S. J. Khairnar, A.R. Bendale, Chalcone scaffolds, bioprecursors of flavonoids: chemistry, bioactivities, and pharmacokinetics, *Molecules* 26 (2021) 7177, <https://doi.org/10.3390/molecules26237177>.
- [75] S.-H. Kim, E. Lee, K.H. Baek, H.B. Kwon, H. Woo, E.-S. Lee, Y. Kwon, Y. Na, Chalcones, inhibitors for topoisomerase I and cathepsin B and L, as potential anti-cancer agents, *Bioorg. Med. Chem. Lett.* 23 (2013) 3320–3324, <https://doi.org/10.1016/j.bmcl.2013.03.106>.
- [76] N. Raghav, M. Singh, SAR studies of differently functionalized chalcones based hydrazones and their cyclized derivatives as inhibitors of mammalian cathepsin B and cathepsin H, *Bioorg. Med. Chem.* 22 (2014) 4233–4245, <https://doi.org/10.1016/j.bmc.2014.05.032>.
- [77] I. Ravish, N. Raghav, SAR studies of differently functionalized 4'-phenylchalcone based compounds as inhibitors of cathepsins BH and L, *RSC Adv.* 5 (2015) 50440–50453, <https://doi.org/10.1039/C5RA00357A>.
- [78] N. Raghav, S. Garg, N. formylpyrazolines and N benzylpyrazolines as novel inhibitors of mammalian cathepsin B and cathepsin H, *Bioorg. Chem.* 57 (2014) 43–50, <https://doi.org/10.1016/j.bioorg.2014.07.012>.
- [79] N. Raghav, S. Garg, SAR studies of o-hydroxychalcones and their cyclized analogs and study them as novel inhibitors of cathepsin B and cathepsin H, *Eur. J. Pharm. Sci.* 60 (2014) 55–63, <https://doi.org/10.1016/j.ejps.2014.04.006>.
- [80] N. Raghav, R. Kaur, Chalcones, semicarbazones and pyrazolines as inhibitors of cathepsins B, H and L, *Int. J. Biol. Macromol.* 80 (2015) 710–724, <https://doi.org/10.1016/j.ijbiomac.2015.07.029>.
- [81] K.-H. Jeon, F. Lee, K.-Y. Jun, J.-E. Eom, S.Y. Kwak, Y. Na, Y. Kwon, Neuroprotective effect of synthetic chalcone derivatives as competitive dual inhibitors against μ -calpain and cathepsin B through the downregulation of tau phosphorylation and insoluble A β peptide formation, *Eur. J. Med. Chem.* 121 (2016) 433–444, <https://doi.org/10.1016/j.ejmech.2016.06.008>.
- [82] D.H. Soliman, A.M. Farrag, O. Omran, Design, synthesis and in-silico studies of novel chalcones as anti-prostate cancer and cathepsin B inhibitors, *J. Appl. Pharm. Sci.* 7 (2017) 010–020, <https://doi.org/10.7324/JAPS.2017.70702>.
- [83] R. Kaur, N. Raghav, Some thiocarbamoyl based novel anticathepsin agents, *Bioorg. Chem.* 104 (2020), 104174, <https://doi.org/10.1016/j.bioorg.2020.104174>.
- [84] G. Sul, T. Li, B. Zhang, R. Wang, H. Hao, W. Zhou, Recent advances on synthesis and biological activities of aurones, *Bioorg. Med. Chem.* 29 (2021), 115895, <https://doi.org/10.1016/j.bmc.2020.115895>.
- [85] B. Saroha, G. Kumar, E. Lathwal, S. Kumar, M. Kumari, N. Mor, N. Raghav, S. Kumar, Synthesis of propynyloxy substituted some novel aurones as potent cathepsin B inhibitors, *Chem. Data Collect.* 31 (2021), 100630, <https://doi.org/10.1016/j.cdc.2020.100630>.
- [86] B. Saroha, G. Kumar, S. Kumar, M. Kumari, M. Rani, N. Raghav, P.K. Sahoo, S. Ghosh, S. Mahata, V.D. Nasare, Ultrasound assisted a one pot multicomponent and greener synthesis of 1, 2, 3-triazole incorporated aurone hybrids: cathepsin B inhibition, anti-cancer activity against AGS cell line, and in-silico docking evaluation, *Curr. Res. Green. Sustain. Chem.* 5 (2022), 100295, <https://doi.org/10.1016/j.crgsc.2022.100295>.
- [87] B. Saroha, G. Kumar, S. Kumar, M. Kumari, M. Rani, N. Raghav, P.K. Sahoo, S. Ghosh, S. Mahata, V.D. Nasare, Novel 1,2,3-triazole-aurone hybrids as cathepsin B inhibitors: one-pot synthesis, anti-proliferative, and drug modeling studies, *Eur. J. Med. Chem. Rep.* (2022), 100056, <https://doi.org/10.1016/j.ejmech.2022.100056>.
- [88] M.T.M. Sarg, S.S. El-Shaer, Efficient utilization of 6-aminouracil to synthesize fused and related heterocyclic compounds and their evaluation as prostate cytotoxic agents with cathepsin B inhibition, *Open J. Med. Chem.* 4 (2014) 22, <https://doi.org/10.4236/ojmc.2014.42003>.
- [89] N. Raghav, M. Singh, Acyl hydrazides and triazoles as novel inhibitors of mammalian cathepsin B and cathepsin H, *Eur. J. Med. Chem.* 77 (2014) 231–242, <https://doi.org/10.1016/j.ejmech.2014.03.007>.
- [90] N. Raghav, R. Kaur, Synthesis and evaluation of some semicarbazone- and thiosemicarbazone-based cathepsin B inhibitors, *Med. Chem. Res.* 23 (2014) 4669–4679, <https://doi.org/10.1007/s00044-014-1036-7>.
- [91] N. Raghav, S. Jangra, A. Kumar, S. Bhattacharya, Quinazoline derivatives as cathepsins B, H and L inhibitors and cell proliferating agents, *Int. J. Biol. Macromol.* 94 (2017) 719–727, <https://doi.org/10.1016/j.ijbiomac.2016.10.001>.
- [92] N. Raghav, M. Singh, SAR studies of some acetophenone phenylhydrazone based pyrazole derivatives as anticathepsin agents, *Bioorg. Chem.* 75 (2017) 38–49, <https://doi.org/10.1016/j.bioorg.2017.08.006>.
- [93] S.F.P. Braga, L.C. Martins, E.B. da Silva, P.A. Sales Júnior, S.M.F. Murta, A. J. Romanha, W.T. Soh, H. Brandstetter, R.S. Ferreira, R.B. de Oliveira, Synthesis and biological evaluation of potential inhibitors of the cysteine proteases cruzain and rhodesain designed by molecular simplification, *Bioorg. Med. Chem.* 25 (2017) 1889–1900, <https://doi.org/10.1016/j.bmc.2017.02.009>.
- [94] B. Luklić, U. Grošelj, M. Novinec, J. Svete, Synthesis of novel 5-(N-boc-N-benzyl-2-aminoethyl)-7-oxo-4, 7-dihydropyrazolo [1, 5-a] pyrimidin-3-carboxamides and their inhibition of cathepsins B and H, *Acta Chim. Slov.* 64 (2017) 782–789, <https://doi.org/10.17344/acsl.2017.3421>.
- [95] M. Rischer, L. Raguz, H. Guo, F. Kelf, G. Diekert, T. Goris, C. Beemelmans, Biosynthesis, synthesis, and activities of barnesin A, a NRPS-PKS hybrid produced by an anaerobic epsilonproteobacterium, *ACS Chem. Biol.* 13 (2018) 1990–1995, <https://doi.org/10.1021/acscchembio.8b00445>.
- [96] S. Garg, N. Raghav, 2,5-diaryloxadiazoles and their precursors as novel inhibitors of cathepsins B, H and L, *Bioorg. Chem.* 67 (2016) 64–74, <https://doi.org/10.1016/j.bioorg.2016.05.003>.
- [97] A. Mitrović, B. Mirković, I. Sostić, S. Gobec, J. Kos, Inhibition of endopeptidase and exopeptidase activity of cathepsin B impairs extracellular matrix degradation and tumor invasion, *Biol. Chem.* 397 (2016) 165–176, <https://doi.org/10.1515/bch-2015-0236>.

NEW International Journal Of Biological Macromolecules

Overview

Impact Factor

6.953




H Index

144



Impact Factor

7.713



I. Basic Journal Info

Country

Netherlands (<https://www.scijournal.org/country/netherlands>)

Journal ISSN: 01418130, 18790003

Publisher: Elsevier BV (<https://www.scijournal.org/publishers/elsevier-bv>)

History: 1979-ongoing

Journal Homepage: Link

(http://www.elsevier.com/wps/find/journaldescription.cws_home/525446/description#description)